



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

# The role of the systematic inflammatory response in predicting outcomes in patients with advanced inoperable cancer: systematic review and meta analysis

### Citation for published version:

Dolan, R, McSorley, S, Horgan, PG, Laird, B & McMillan, DC 2017, 'The role of the systematic inflammatory response in predicting outcomes in patients with advanced inoperable cancer: systematic review and meta analysis', *Critical Reviews in Oncology/Hematology*. <https://doi.org/10.1016/j.critrevonc.2017.06.002>

### Digital Object Identifier (DOI):

[10.1016/j.critrevonc.2017.06.002](https://doi.org/10.1016/j.critrevonc.2017.06.002)

### Link:

[Link to publication record in Edinburgh Research Explorer](#)

### Document Version:

Peer reviewed version

### Published In:

Critical Reviews in Oncology/Hematology

### General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## **Abstract:**

**Introduction:** Cancer remains a leading cause of death worldwide. While a curative intent is the aim of any surgical treatment many patients either present with or go on to develop disseminated disease requiring systemic anti-cancer therapy with a palliative intent. Given their limited life expectancy appropriate allocation of treatment is vital. It is recognised that systemic chemoradiotherapy may shorten the quality/quantity of life in patients with advanced cancer. It is against this background that the present systematic review and meta-analysis of the prognostic value of markers of the systemic inflammatory response in patients with advanced cancer was conducted.

**Methods:** An extensive literature review using targeted medical subject headings was carried out in the MEDLINE, EMBASE, and CDSR databases until the end of 2016. Titles were examined for relevance and studies relating to duplicate datasets, that were not published in English and that did not have full text availability were excluded. Full texts of relevant articles were obtained and were then examined to identify any further relevant articles.

**Results:** The majority of studies were retrospective. The systemic inflammatory response, as evidenced by a number of markers at clinical thresholds, was reported to have independent prognostic value, across tumour types and geographical locations. In particular, C-reactive protein (CRP, 63 studies), albumin (33 studies) the Glasgow Prognostic Score (GPS, 44 studies) and the Neutrophil Lymphocyte Ratio (NLR, 59 articles) were consistently validated across tumour types and geographical locations. There was considerable variation in the thresholds reported to have prognostic value when CRP and albumin were examined. There was less variation in the thresholds reported for NLR and still less for the GPS.

**Discussion:** The systemic inflammatory response, especially as evidenced by the GPS and NLR, has reliable prognostic value in patients with advanced cancer. Further prospective studies of their clinical utility in randomised clinical trials and in treatment allocation are warranted.

## Introduction:

Cancer remains one of the leading cause of death worldwide and is responsible for 7.6 million deaths per year. Therefore, while a curative intent is the aim of any surgical treatment many patients either present with or go onto develop disseminated disease requiring systemic anti-cancer therapy with a palliative intent. Given that patients with advanced cancer have a limited life expectancy appropriate treatment selection becomes vital. Indeed, the paradigm of precision medicine (right treatment, right patient, right time) is in the vanguard of oncology treatment, and if applied outcomes for all patients would improve irrespective of new treatment availability.<sup>1</sup>

However, optimal allocation of treatment remains elusive. There is increasing evidence that inappropriate anti-cancer treatment does not improve quality of life or survival<sup>2,5</sup>. A National Clinical Enquiry into Patient Outcome and Death (NCEPOD) reported that chemotherapy hastened or directly caused the death of over 25% of patients who died within 30 days of receiving treatment<sup>4</sup>. This need for caution has been further illustrated by a randomised control trial comparing early palliative and standard oncological care in patients with metastatic non-small cell lung cancer conducted by Temel *et al*<sup>5</sup>. In this randomised trial patients who received palliative care early not only maintained better quality of life scores but also had a significantly longer median survival<sup>5</sup>. These reports provide a persuasive argument for optimising the stratification of anti-cancer therapy in patients with advanced cancer. Therefore, it is important to examine the criteria that may be used to effectively stratify patients as to their likely survival prior to the allocation of treatment in patients with advanced cancer.

In the setting of patients with advanced cancer, Tumour, Node, Metastasis (TNM) staging has little discriminatory prognostic value and other patient related measures such as weight loss, performance status and quality of life have superior prognostic value. Therefore, the decision to

proceed with systemic therapy is frequently based on these parameters by an oncologist and primarily on the basis of subjective clinical observation. More recently, measurement of skeletal muscle mass made from CT scans has been proposed to be useful in this context<sup>6</sup>. Nevertheless, it is clear that the potential for sub-optimal allocation of anti-cancer therapy is considerable.

Recently, in a systematic review of prognostic tools in patients with advanced cancer, it was reported that a number of prognostic tools had been validated in different centres<sup>7</sup>. It was striking that the majority of these validated tools were based on subjective criteria, in particular the assessment of physical function. Only one validated prognostic tool the GPS (Glasgow Prognostic Score), assessing the magnitude of the systemic inflammatory response, was based exclusively on objective criteria. Indeed, there is now strong evidence that the chronic systemic inflammatory response results in classical features of cancer cachexia, including the preferential loss of lean muscle mass<sup>8-10</sup>. Indeed, studies have shown a direct relationship between systemic inflammation measured by the GPS and NLR (Neutrophil Lymphocyte Ratio) and elevation of inflammatory cytokines, adipokines and other biochemical disturbances associated with loss of lean muscle mass and reduced performance status<sup>8,11-14</sup>. Recently, Laird and co-workers showed that in a large cohort study in two international bio banks, the combination of performance status and the systemic inflammatory response (SIR) as measured by the mGPS (modified Glasgow Prognostic Score) improved the prediction of outcomes of patients with advanced cancer<sup>15</sup>. Furthermore, they showed that quality of life was independently associated with both performance and the GPS<sup>16</sup>.

Therefore, from the above and with the introduction of immunotherapeutic agents for advanced inoperable cancer it is timely to review the role of the markers of systemic inflammatory response in predicting outcomes in patients with advanced inoperable cancer.

## **Methods:**

The present systematic review and meta-analysis of published literature was undertaken according to a pre-defined protocol described in the PRISMA-P statement. The primary outcome was to assess the prognostic value of the SIR in patients with advanced inoperable cancer treated with chemotherapy, immunotherapy, radiotherapy, best supportive care or a combination of these treatment strategies.

This was carried out by a wide-ranging literature search to identify studies. Medical subject heading (MeSH) terms (Advanced Cancer, CRP, C-Reactive Protein, Albumin, White Cell Count, Neutrophil Count, Lymphocyte Count, Monocyte Count, Platelet Count, Red Blood Cell Count), were used in the US National Library of Medicine (MEDLINE), the Excerpta Medica database (EMBASE) and the Cochrane Database of Systematic Reviews (CDSR) to identify articles.

On completion of the online search, the title and abstract of each identified study was examined for relevance. Studies relating to duplicate datasets, studies not available in English and those published in abstract form only were excluded. Full texts were obtained for all studies deemed potentially relevant. Once further exclusions outlined below were carried out the bibliographies of all included articles were subsequently hand searched to identify any additional studies.

Only articles that reported survival analysis and gave hazard ratios or odds ratios with associated confidence intervals were included in the review. Studies with patients who had failed resections and patients who underwent palliative symptom control procedures were also included.

## **Statistics**

The HRs and 95 % CIs were directly retrieved from the article. If several estimates were reported for the same marker, the multivariate estimate was used in preference to the univariate analysis.

Interstudy heterogeneity among included studies was evaluated by  $I^2$  statistics using the random-effects (DerSimonian – Laird method) model. All P values were 2-sided and  $P < 0.05$  were considered statistical significant. Evidence of publication bias was evaluated using visual inspection of funnel plots. All analyses were performed using Review Manager (RevMan) [Computer program]. Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

## Results

### *Study selection process*

Initial search strategy identified 9546 articles whose titles and abstracts were reviewed (Figure 1). Articles were excluded if initial curative surgery formed part of the treatment regimen (n=3114), where survival was not the primary outcome measure (n=1225), full articles were not available (n=1195), articles examining response to bacterial and viral infection (n=924), articles not carried out in humans (n=2021), articles not published in English (n=219), and those that were a systematic review/meta-analysis (n=149).

This led to a review of the full text of 699 articles. Further articles were excluded if surgery was part of the treatment regimen being examined (n=421), progression free survival (PFS) was the only outcome measured (n=62) and if survival was not expressed as HR (95%CI; n=47). The remaining 169 articles, had their bibliographies reviewed in a systematic manner and this identified a further 29 articles to be included in the final analysis leading to a final total of 198 articles.

### **Figure 1:**

#### *Studies of the prognostic value of C-reactive protein (CRP) in patients with advanced cancer:*

Sixty-three articles with both overall survival (OS) and/or cancer specific survival (CSS) as their primary outcome measures were identified comprising data on 13,498 patients (8,466 deaths) (Supplementary Table). Fifty-four studies were carried out in a retrospective manner while eight were prospective with one study having both prospective and retrospective arms (Supplementary Table). Fifty-four studies used multivariate and nine used univariate survival analysis (Supplementary Table). On meta-analysis of the 55 retrospective studies including 11,761 patients (7,316 deaths) there was a significant association between elevated CRP and survival (HR: 1.97 95%CI 1.76-2.21,  $p<0.00001$ ) with a considerable degree of heterogeneity ( $I^2=92\%$ ). On meta-



analysis of the 9 prospective studies including 1,598 patients (1,009 deaths) there was a significant association between elevated CRP and survival (HR: 1.72 95%CI 1.31-2.26,  $p<0.00001$ ) with a considerable degree of heterogeneity ( $I^2=88\%$ ).

Fifty-six studies examined the relationship with overall survival including 11,787 patients (7,477 deaths), as the primary outcome measure. On meta-analysis, there was a significant association between CRP and overall survival (HR 1.47 95%CI 1.40-1.54,  $p<0.00001$ ) with a considerable degree of heterogeneity ( $I^2=90\%$ , Figure 2). There was variation in the threshold of CRP used in the studies, the most common being  $>10$  mg/L ( $n=19$ ) followed by  $>5$  mg/L ( $n=5$ ). Other thresholds ( $n=32$ ) were used in  $<5$  studies and therefore meta-analysis was not carried out.

On meta-analysis those studies with a threshold of  $>10$ mg/L ( $n=19$ ), including 3,883 patients (3,458 deaths), there was a significant association between CRP and overall survival (HR: 1.73 95%CI 1.55-1.93,  $p<0.00001$ ) with a moderate degree of heterogeneity ( $I^2 = 35\%$ ). These included studies on cancer of the pancreas ( $n=6$ ), lung ( $n=5$ ), lymphoma ( $n=2$ ), HCC ( $n=1$ ), osteosarcoma ( $n=1$ ), prostate ( $n=1$ ), oesophagus ( $n=1$ ), multiple cancers ( $n=1$ ) and renal cells ( $n=1$ ).

On meta-analysis of those studies with a threshold of  $>10$ mg/L and pancreatic cancer ( $n=6$ ) 1,510 patients (1,446 deaths) there was a significant association between CRP and overall survival (HR: 1.64 95%CI 1.28-2.10,  $p<0.0001$ ) with substantial heterogeneity ( $I^2=73\%$ ). In these six studies, there was a variation in their geographical locations including Japan ( $n=2$ ), Korea ( $n=2$ ), Germany ( $n=1$ ) and Australia ( $n=1$ ). The proportion of patients who had a CRP level  $>10$ mg/L with pancreatic cancer was 90% in Japan, 65% in Korea, 63% in Australia and 19% in Germany.

On meta-analysis of those studies with a threshold of  $>10$ mg/L and lung cancer ( $n=5$ ) including 996 patients (960 deaths) there was a significant association between CRP and overall survival (HR: 1.58 95%CI 1.37-1.84,  $p<0.00001$ ) with no heterogeneity ( $I^2=0\%$ ). In these 5 studies, there was a wide variation in their geographical locations including the Czech Rep ( $n=1$ ), UK ( $n=1$ ), Sweden ( $n=1$ ), China ( $n=1$ ) and Japan ( $n=1$ ). The proportion of patients who had a CRP level  $>10$ mg/L and lung cancer was 98% in the Czech Rep, 80% in the UK, 71% in Sweden, 43% in China and 33% in

Japan. Remaining cancer types and geographical locations had <5 studies therefore further meta-analysis was not carried out.

On meta-analysis those studies with a threshold of >5mg/L (n=5), including 961 patients (515 deaths), there was a significant association between CRP and overall survival (HR: 1.66 95%CI 1.15-2.38, p=0.007) with a substantial degree of heterogeneity ( $I^2 = 83\%$ ). These included studies on cancer of the pancreas (n=2), prostate (n=1), renal cells (n=1) and colorectal (n=1). These included studies carried out in Japan (n=3), Belgium (n=1) and Sweden (n=1). The proportion of patients who had a CRP>5mg/L was 100% in Sweden, 66% in Belgium and 50% in Japan. Remaining cancer types and geographical locations had <5 studies therefore further meta-analysis was not carried out.

Ten studies examined the relationship with cancer specific survival including 1711 patients (989 deaths), as its primary outcome measure. On meta-analysis, there was a significant association between CRP and cancer specific survival (HR 2.93 95%CI 2.14-4.01, p<0.00001) with a substantial degree of heterogeneity ( $I^2=66\%$ ). The most common thresholds used on the CSS group were >10 mg/L (n=4) including cancer of the prostate (n=1), breast (n=1), renal cells (n=1) and urothelial (n=1). All thresholds had <5 studies and therefore meta-analysis was not carried out. In the >10mg/L group studies were carried out in the UK (n=3) and Italy (n=1). The proportion of patients who had a CRP level >10mg/L was 64% in the UK and 50% in Italy.

#### *Studies of the prognostic value of albumin (Alb) in patients with advanced cancer:*

Thirty-three articles with both OS (n=29) and/or CSS (n=5) as their primary outcome measures were identified comprising data on 10,288 patients (8,740 deaths) (Supplementary Table). Twenty-eight studies were conducted in a retrospective manner while five were prospective. Twenty-nine articles used multivariate and four univariate survival analysis (Supplementary Table).

Thirty-one studies examined the relationship with overall survival including 9,753 patients (8,493 deaths), as its primary outcome measure. On meta-analysis, there was a significant association between low albumin and overall survival (HR 1.77 95%CI 1.54-2.03,  $p < 0.00001$ ) with a considerable degree of heterogeneity ( $I^2 = 84\%$ , Figure 3). There was variation in the threshold of albumin examined. The most common thresholds examined were  $<35\text{g/L}$  ( $n=13$ ) and  $<30\text{ mg/L}$  ( $n=5$ ). Other thresholds were used in  $<5$  studies ( $n=15$ ) and therefore meta-analysis was not carried out.

On meta-analysis those studies with a threshold of  $<35\text{g/L}$  ( $n=13$ ), including 2,127 patients (1,831 deaths), there was a significant association between low albumin and overall survival (HR: 2.21 95%CI 1.60-3.06,  $p < 0.00001$ ) with a considerable degree of heterogeneity ( $I^2 = 79\%$ ). These included studies on cancer of the pancreas ( $n=5$ ), biliary tract ( $n=2$ ), multi anatomical sites ( $n=1$ ), breast ( $n=1$ ), lung ( $n=1$ ), HCC ( $n=1$ ), colorectal ( $n=1$ ) and multiple myeloma ( $n=1$ ). These included studies carried out in Korea ( $n=6$ ), Japan ( $n=3$ ), Singapore ( $n=1$ ), Canada ( $n=1$ ), Belgium ( $n=1$ ), France ( $n=1$ ), Spain ( $n=1$ ), Australia ( $n=1$ ), and the UK ( $n=1$ ). The proportion of patients who had an albumin  $<35\text{g/L}$  was 51% in Korea, 49% in Spain, 31% in Belgium, 26% in the UK and 16% in France.

On meta-analysis of those studies with a threshold of  $<35\text{g/L}$  and pancreatic cancer ( $n=5$ ) 910 patients (834 deaths) there was a significant association between reduced albumin and overall survival (HR: 1.96 95%CI 1.04-3.69,  $p=0.04$ ) with substantial heterogeneity ( $I^2=85\%$ ). In these five studies, there was a variation in their geographical locations including Korea ( $n=2$ ), Japan ( $n=1$ ), Australia ( $n=1$ ) and Belgium ( $n=1$ ). The proportion of patients who had an albumin level  $<35\text{g/L}$  with pancreatic cancer was 31% in Belgium and 42% in Australia.

On meta-analysis of those studies with a threshold of  $<30\text{g/L}$  ( $n=5$ ), including 1,319 patients (1,192 deaths), there was a significant association between low albumin and overall survival (HR: 1.57 95%CI 1.26-1.95,  $p < 0.0001$ ) with a minimal degree of heterogeneity ( $I^2 = 14\%$ ). These included studies on cancer of the lung ( $n=2$ ), gastric ( $n=1$ ), renal cells ( $n=1$ ), and multiple

anatomical sites (n=1). These included studies carried out in the US (n=1), Taiwan (n=1), Japan (n=1), Turkey (n=1) and Sweden (n=1). The proportion of patients who had an albumin <30g/L was 49% on Taiwan, 39% in the Japan, 20% in Turkey and 17% in Sweden.

*Studies of the prognostic value of white cell count (WCC) in patients with advanced cancer:*

Four articles with both OS (n=3) and/or CSS (n=1) as their primary outcome measures were identified comprising data on 1,593 patients (1,440 deaths) (Supplementary Table). All four were retrospective multivariate survival studies carried out in cancer of the lung (n=2), renal cells (n=1) and multiple anatomical sites (n=1). There was variation in the level of WCC used between different papers including  $>10 \times 10^9/L$  (n=2),  $>10.2 \times 10^9/L$  for males and  $>10.6 \times 10^9/L$  for females (n=1), and  $>11 \times 10^9/L$  for both sexes (n=1). Geographically studies were carried out in the UK (n=2), US (n=1) and Italy (n=1). The proportion of patients who had an elevated WCC was 24% in the US, 28% in the UK and 28% in Italy. Due to the small number of studies, meta-analysis was not carried out.

*Studies of the prognostic value of neutrophils in patients with advanced cancer:*

Nine articles with both OS (n=7) and/or CSS (n=2) as their primary outcome measures were identified comprising data on 2,870 patients (2,266 deaths) (Supplementary Table). Seven studies were conducted in a retrospective manner while two were prospective. (Supplementary Table). Five articles reported significance on multivariate and two articles reported significance on univariate survival analysis. There was variation in the levels of neutrophils used in individual papers including neutrophil count  $\geq$  upper limit of normal (ULN) without defining it explicitly (n=3), neutrophil count  $>7.5 \times 10^9$  cells/ml (n=1), neutrophil count  $>3.41 \times 10^9$  cells/ml (n=1),

absolute neutrophil count (ANL)  $>4.7 \times 10^9/L$  (n=1),  $ANC \geq 7500$  (n=1), log of readings above normal which was defined as  $>7 \times 10^9/L$  (n=1) and  $>8 \times 10^9/L$  (n=1), .

Seven studies examined the relationship with overall survival including 2,364 patients (1,999 deaths), as its primary outcome measure. On meta-analysis, there was a significant association between elevated neutrophils and overall survival (HR 1.89 95%CI 1.25-2.85,  $p=0.002$ ) with a considerable degree of heterogeneity ( $I^2=87\%$ ). Studies were in melanoma (n=2), renal (n=1), lung (n=1), breast (n=1), mesothelioma (n=1) and lung (n=1) cancer. Geographically studies were carried out in France (n=2) and Italy (n=2), USA (n=1), China (n=1) and Australia (n=1). The proportion of patients who had elevated Neutrophils was 32% in Australia, 28% in France, 19% in the USA and 12% in Italy.

Two studies examined the relationship with cancer specific survival including 506 patients (267 deaths), as its primary outcome measure. Due to the small number of studies, meta-analysis was not carried out.

*Studies of the prognostic value of lymphocytes in patients with advanced cancer:*

Eleven articles with OS as their primary outcome measures were identified comprising data on 2,517 patients (2,148 deaths) (Supplementary Table). Ten studies were conducted in a retrospective manner and one prospectively. Nine studies reported significance on multivariate survival analysis and two on univariate survival analysis. (Supplementary Table). On meta-analysis, there was a significant association between lower lymphocyte levels and overall survival (HR 1.68 95%CI 1.35-2.09,  $p<0.00001$ ) with a substantial degree of heterogeneity ( $I^2=68\%$ ).

There was considerable variation in the lymphocyte thresholds used in each study including continuous readings (n=1),  $<0.5 \times 10^9/L$  (n=1),  $<0.7 \times 10^9/L$  (n=1),  $>2 \times 10^9/L$  (n=2),  $<1 \times 10^9/L$  (n=2),  $\geq 0.45 \times 10^9/L$  (n=1),  $<2.25 \times 10^9/L$  (n=1),  $<1.4 \times 1 \times 10^9/L$  (n=1), and  $2.70 \times 10^9/L$  (n=1). These included

studies on cancer of the pancreas (n=3), lymphoma (n=1), lung (n=1), nasopharyngeal (n=1), mesothelioma (n=1), colorectal (n=1), cervical (n=1), melanoma (n=1) and multiple cancer types (n=1). Geographically studies were carried out in China (n=3), US (n=3), France (n=2), Japan (n=2) and Korea (n=1). The proportion of patients who had low lymphocytes was 75% in Korea, 48% in US, 47% in China, 45% in Japan and 32% in France. All eleven studies used chemotherapy as the treatment modality. No specific lymphocyte thresholds had more than four studies and therefore no further meta-analysis was carried out.

*Studies of the prognostic value of monocytes in patients with advanced cancer:*

Five articles with OS as their primary outcome measures were identified comprising data on 1,367 patients (1,152 deaths) (Supplementary Table). All five studies were conducted in a retrospective multivariate manner, used chemotherapy as the treatment regime of choice and conducted their analysis in a multivariate manner. On meta-analysis of there was a significant association between elevated monocytes and survival (HR: 1.40 95%CI 1.05-1.87, p=0.02) with a substantial degree of heterogeneity ( $I^2=66\%$ ). There was considerable variation in the levels of monocytes used including  $>0.8 \times 10^9/L$  (n=1),  $\geq 0.64 \times 10^9/L$  (n=1),  $\geq 0.45 \times 10^9/L$  (n=1),  $\geq 0.35 \times 10^9/L$  (n=1) and  $\geq 0.55 \times 10^9/L$  (n=1). There was also variation in the types of cancer examined including lung (n=2), lymphoma (n=1), nasopharyngeal (n=1) and colorectal metastasis (n=1). In terms of geographical locations, the studies were carried out in China (n=3), Korea (n=1) and Italy (n=1). The proportion of patients who had high monocytes was 57% in China, 50% in Korea, and 23% in Italy. No specific monocyte thresholds had more than four studies and therefore no further meta-analysis was carried out.

*Studies of the prognostic value of platelets in patients with advanced cancer:*

Eight articles with both OS (n=7) and/or CSS (n=1) as their primary outcome measures were identified comprising data on 4,850 patients (2,422 deaths) (Supplementary Table). Seven studies were conducted in a retrospective manner while one was prospective (Supplementary Table). All eight articles reported multivariate survival analysis.

Seven studies examined the relationship with overall survival including 4,653 patients (2,293 deaths), as its primary outcome measure. On meta-analysis of there was a significant association between elevated platelets and survival (HR: 1.47 95%CI 1.12-1.93, p=0.006) with a considerable degree of heterogeneity ( $I^2=92\%$ ). There was variation in the thresholds of platelets examined including a platelet count  $>300 \times 10^9 /L$  (n=1),  $>360 \times 10^9 /L$  (n=1),  $<130 \text{ g/L}$  (n=1),  $>350 \times 10^9 /L$  (n=1),  $>450 \times 10^9 /L$  (n=1),  $\geq \text{ULN}$  (n=1) and continuous readings (n=1). There was also variation in the type of cancers being examined including lung (n=1), oropharyngeal (n=1), pleural mesothelioma (n=1), nasopharyngeal (n=1), pancreatic (n=1), renal (n=1) and multiple cancers (n=1). Geographically studies were carried out in US (n=3), China (n=2), France (n=1) and Sweden (n=1). The proportion of patients who had elevated platelet counts was 30% in Sweden, 24% in the US, 15% in China and 11% in France. However, no specific platelet thresholds had more than four studies and therefore no further meta-analysis was carried out.

*Studies of the prognostic value of the Glasgow Prognostic Score (GPS/mGPS) in patients with advanced cancer:*

Forty-four articles with both OS (n=37) and/or CSS (n=9) as their primary outcome measures were identified comprising data on 12,578 patients (10,745 deaths) (Supplementary Table). Thirty-two studies were conducted in a retrospective manner while twelve were prospective

(Supplementary Table). Forty studies reported multivariate and four reported univariate survival analysis (Supplementary Table). On meta-analysis of the 32 retrospective studies including 9,472 patients (7,936 deaths) there was a significant association between elevated GPS/mGPS and survival (HR: 1.93 95%CI 1.76-2.13,  $p < 0.00001$ ) with a moderate degree of heterogeneity ( $I^2 = 42\%$ ). On meta-analysis of the 12 prospective studies including 3,244 patients (2,809 deaths) there was a significant association between elevated GPS/mGPS and survival (HR: 2.09 95%CI 1.69-2.57,  $p = 0.0001$ ) with a substantial degree of heterogeneity ( $I^2 = 69\%$ ).

Thirty-six studies examined the relationship with overall survival including 11,441 patients (10,022 deaths), as its primary outcome measure. On meta-analysis, there was a significant association between GPS and overall survival (HR 2.06 95%CI 1.86-2.28,  $p < 0.00001$ ) with a substantial degree of heterogeneity ( $I^2 = 56\%$ , Figure 4). These included studies on cancer of multiple anatomical sites ( $n = 7$ ), gastric ( $n = 7$ ), lung ( $n = 5$ ), pancreas ( $n = 5$ ), colon ( $n = 3$ ), lymphoma ( $n = 1$ ), biliary tract ( $n = 1$ ), bladder ( $n = 1$ ), haematological ( $n = 1$ ), prostate ( $n = 1$ ), renal cell ( $n = 1$ ), oesophagus ( $n = 1$ ), HCC ( $n = 1$ ) and cervix ( $n = 1$ ).

On meta-analysis those studies carried out in multiple anatomical sites ( $n = 7$ ), including 5,804 patients (5,139 deaths), there was a significant association between elevated GPS/mGPS and overall survival (HR: 2.22 95%CI 1.81-2.71,  $p < 0.00001$ ) with a moderate degree of heterogeneity ( $I^2 = 65\%$ ). These included studies carried out in the UK ( $n = 2$ ), Australia ( $n = 2$ ), Japan ( $n = 1$ ), Norway ( $n = 1$ ) and Brazil ( $n = 1$ ). The proportion of patients who had an elevated GPS was 93% in Japan, 77% in the UK, 69% in Norway, 46% in Australia and 20% in Brazil.

On meta-analysis those studies carried out in gastric cancer ( $n = 7$ ), including 1,283 patients (5139 deaths), there was a significant association between elevated GPS/mGPS and overall survival (HR: 2.08 95%CI 1.58-2.74,  $p < 0.00001$ ) with a moderate degree of heterogeneity ( $I^2 = 40\%$ ). These included studies carried out in the Japan ( $n = 2$ ), Korea ( $n = 2$ ), Taiwan ( $n = 1$ ), UK ( $n = 1$ ) and Czech



Rep (n=1). The proportion of patients who had an elevated GPS was 74% in Taiwan, 73% in the UK, 52% in the Czech Rep, 49% in Japan and 42% in Korea.

On meta-analysis those studies carried out in lung cancer (n=5), including 1,104 patients (708 deaths), there was a significant association between elevated GPS and overall survival (HR: 2.05 95%CI 1.52-2.77,  $p < 0.00001$ ) with a substantial degree of heterogeneity ( $I^2 = 55\%$ ). These included studies carried out in the UK (n=2), China (n=2) and Greece (n=1). The proportion of patients who had an elevated GPS was 76% in the UK, 33% in China and 29% in Greece.

On meta-analysis those studies carried out in pancreatic cancer (n=5), including 735 patients (719 deaths), there was a significant association between elevated GPS and overall survival (HR: 1.91 95%CI 1.29-2.83,  $p = 0.001$ ) with a substantial degree of heterogeneity ( $I^2 = 70\%$ ). These included studies carried out in the Japan (n=3), Australia (n=1) and the UK (n=1). The proportion of patients who had an elevated GPS was 70% in the UK, 63% in Australia and 36% in Japan.

Nine studies examined cancer specific survival including 1,137 patients (723 deaths), as its primary outcome measure. On meta-analysis, there was a significant association between elevated GPS and cancer specific survival (HR 1.69 95%CI 1.48-1.92,  $p < 0.00001$ ) with a minimal degree of heterogeneity ( $I^2 = 4\%$ ). These included studies on cancer of the colon (n=3), lung (n=2), gastro-oesophageal (n=2), breast (n=1) and renal cells (n=1). These included studies carried out in the UK (n=5), Japan (n=2) and China (n=2). The proportion of patients who had an elevated GPS was 77% in China, 65% in the UK and 43% in Japan. However, since no cancer type or country had more than four studies further meta-analysis was not carried out.

*Studies of the prognostic value of neutrophil lymphocyte ratio (NLR) in patients with advanced cancer:*

Fifty-nine articles with both OS (n=58) and/or CSS (n=2) as their primary outcome measures were identified comprising data on 16,921 patients (12,801 deaths) (Supplementary Table). Forty-three of these were conducted in a retrospective manner while sixteen were prospective. Fifty-five studies reported multivariate and four reported univariate survival analysis (Supplementary Table). On meta-analysis of the 43 retrospective studies including 10,870 patients (8,044 deaths) there was a significant association between elevated NLR and survival (HR: 1.78 95%CI 1.59-1.98,  $p<0.00001$ ) with a considerable degree of heterogeneity ( $I^2=77\%$ ). On meta-analysis of the 16 prospective studies including 5,898 patients (4,733 deaths) there was a significant association between elevated NLR and survival (HR: 1.63 95%CI 1.41-1.88,  $p<0.00001$ ) with a substantial degree of heterogeneity ( $I^2=67\%$ ).

Fifty-eight studies examined the relationship with overall survival including 16,405 patients (12,675 deaths) as its primary outcome measure. On meta-analysis, there was a significant association between NLR and overall survival (HR 1.71 95%CI 1.57-1.86,  $p<0.00001$ ) with a substantial degree of heterogeneity ( $I^2=79\%$ , Fig 5). The most common NLR thresholds used were  $\geq 5$  (n=19),  $\geq 4$  (n=5) and  $\geq 3$  (n=12). Other thresholds were used in <5 studies and therefore meta-analysis was not carried out (n=23).

On meta-analysis those studies with a threshold of  $\geq 5$  (n=19), including 5,506 patients (4,613 deaths) there was a significant association between elevated NLR and overall survival (HR: 1.64 95%CI 1.42-1.89,  $p<0.00001$ ) with a substantial degree of heterogeneity ( $I^2 = 57\%$ ). These included cancer of the pancreas (n=5), lung (n=4), colorectal (n=3), multiple anatomical sites (n=2), mesothelioma (n=1), prostate (n=2), cholangiocarcinoma (n=1) and HCC (n=1).

On meta-analysis of those studies with a threshold of  $\geq 5$  and pancreatic cancer (n=5) 1009 patients (942 deaths) there was a significant association between an  $NLR \geq 5$  and overall survival (HR: 1.78 95%CI 1.30-2.44,  $p=0.0003$ ) with substantial heterogeneity ( $I^2=56\%$ ). In these five

studies, there was a variation in their geographical locations including Japan (n=2), Australia (n=1), Korea (n=1) and China (n=1). The proportion of patients who had an NLR $\geq$ 5 with pancreatic cancer 48% in Australia, 29% in Korea, and 20% in Japan. No country had more than 4 studies and therefore no further meta-analysis was carried out.

On meta-analysis those studies with a threshold of  $\geq$ 4 (n=5), including 834 patients (588 deaths), there was a significant association between elevated NLR and overall survival (HR: 2.08 95%CI 1.45-3.00, p<0.0001) with a substantial degree of heterogeneity ( $I^2 = 57\%$ ). These included cancer of the lung (n=1), colorectal (n=1), B-cell lymphoma (n=1), T-cell lymphoma (n=1) and gastric (n=1). In these five studies, there was a variation in their geographical locations including Japan (n=2), UK (n=1), Peru (n=1) and Austria (n=1). The proportion of patients who had an NLR $\geq$ 4 was 40% in Japan, 35% in Peru, 32% in the UK and 19% in Austria.

On meta-analysis those studies with a threshold of  $\geq$ 3 (n=12), including 4,195 patients (3,130 deaths), there was a significant association between elevated NLR and overall survival (HR: 1.75 95%CI 1.53-2.01, p<0.00001) with a substantial degree of heterogeneity ( $I^2=56\%$ ). These included cancer of the renal cells (n=3), prostate (n=3), gastric (n=3), melanoma (n=1), colorectal (n=1) and multiple anatomical sites (n=1). These included studies carried out in the Korea (n=2), US/Israel (n=2), China (n=2), Italy (n=2), Australia (n=1), Canada (n=1), Taiwan (n=1) and the UK (n=1). The proportion of patients who had an NLR $\geq$ 3 was 71% in the US/Israel, 53% in Korea, 52% in Australia, 51% in Taiwan, 47% in the UK, 42% in China and 30% in Italy. No tumour site had more than four studies and therefore no further meta-analysis was carried out.

*Studies of the prognostic value of lymphocyte monocyte ratio (LMR) in patients with advanced cancer:*

Eleven articles with both OS (n=11) and/or CSS (n=1) as their primary outcome measures were identified comprising data on 5,043 patients (3,842 deaths) (Supplementary Table). All 11 studies were retrospective and multivariate analysis was carried out. On meta-analysis, there was a significant association between a low LMR and overall survival (HR 1.84 95%CI 1.64-2.07,  $p<0.00001$ ) with minimal heterogeneity ( $I^2=8\%$ , Figure 6). There was a variety of LMR thresholds used in each study including  $\leq 2.6$  (n=1),  $< 2.8$  (n=1),  $\geq 2.475$  (n=1),  $< 2.11$  (n=1),  $> 5.22$  (n=1),  $\leq 4.56$  (n=1),  $\leq 5.07$  (n=1),  $\leq 3.4$  (n=1),  $\leq 2.11$  (n=1),  $\leq 3.11$  (n=1) and low LMR but no figures given (n=1). These included studies on lung cancer (n=2), lymphoma (n=2), nasopharyngeal cancer (n=3) Hodgkin's lymphoma (n=2), and colorectal (n=2). Geographically the studies were carried out in China (n=5), Korea (n=3), Taiwan (n=1), Hungary (n=1) and Italy (n=1). The proportion of patients who had low LMRs was 53% in Italy, 52% in Korea 45% in China and 41% in Taiwan. No specific LMR thresholds had more than four studies and therefore no further meta-analysis was carried out.

*Studies of the prognostic value of platelet lymphocyte ratio (PLR) in patients with advanced cancer:*

Twelve articles with both OS (n=12) and/or CSS (n=2) as their primary outcome measures were identified comprising data on 5,733 patients (2,611 deaths) (Supplementary Table). Ten studies were conducted in a retrospective manner and two prospectively. Eleven studies were also conducted in a multivariate and one in a univariate manner (Supplementary Table). On meta-analysis, there was a significant association between an elevated PLR on overall survival (HR 1.49 95%CI 1.10-1.84,  $p=0.0003$ ) with considerable heterogeneity ( $I^2=82\%$ , Figure 7). There was a variety of PLR thresholds used in each study including  $> 111.23$  (n=1),  $\geq 190$  (n=1),  $> 153.44$  (n=1),  $> 322$  (n=1),  $> 146$  (n=1),  $> 200$  (n=1),  $\geq 152.6$  (n=1),  $\geq 250$  (n=1),  $> 119.50$  (n=1),  $\geq 150$  (n=1),  $> 162$  (n=1) and one study which simply stated elevated PLR without given a numerical value. These included studies on cancer of the lung (n=5), nasopharynx (n=1), cervix (n=1), prostate (n=1), pancreas (n=2), colorectal (n=1) and liver (n=1). Geographically studies were located in China (n=6), Japan (n=2), Turkey (n=1), Austria (n=1), Australia (n=1) and the US (n=1). The proportion

of patients who had an elevated PLR was 61% in Australia, 59% in Japan, 50% in Turkey, 31% in China, 29% in Austria and 20% in the US. No specific PLR thresholds had more than four studies and therefore no further meta-analysis was carried out.

*Studies of the prognostic value of other markers/scores of the systemic inflammatory response in patients with advanced cancer:*

During the course of this review several studies (n=6) were identified which could not be assigned to one of the above groupings (Supplementary Table). Two studies focused on the CRP/Albumin ratio (CAR). The first such study was by Zhou et al<sup>17</sup> from China. In this multivariate survival analysis on patients with small cell lung cancer a CRP/Alb ratio  $\geq 0.441$  was shown to be related to a statistically significant worse OS (HR: 1.34 95%CI 1.04-1.73 p=0.025). The second such study by Yamashita et al<sup>18</sup> from Japan. In this multivariate survival analysis on patients with prostate cancer a CRP/Alb ratio  $\geq 7$  was shown to be related to a statistically non-significant worse overall survival (HR: 2.34 95%CI 0.91-6.05 p=0.08).

Two further studies focused on the relationship between globulin, albumin and survival. Shibutani et al<sup>19</sup> in Japan reported that the albumin/globulin ratio predicted overall survival (HR: 2.247, 95%CI 1.069-4.722, p=0.033) independent of the NLR. Yao et al<sup>20</sup> in China reported that in patients with advanced NSCLC, the globulin/albumin ratio (GAR)  $>0.58$  and an Alb $<35\text{g/L}$  was associated with poorer OS (GAR HR 1.65, 95%CI 1.20-2.26, p=0.002, Alb HR 1.92, 95%CI 1.10-3.36, p=0.022). Chan et al<sup>21</sup> in China reported that, in patients with HCC, the albumin-to-alkaline phosphatase ratio (AAPR)  $>0.68$  predicted poorer OS (HR 2.185, 95%CI, 1.780-2.683, p<0.001).

Finally, Zhou et al<sup>17</sup> in China reported that, in patients with SCLC, the CRP/Globulin ratio  $\geq 1.29$  predicted poorer OS in both the testing (HR 1.35, 95%CI, 1.61-1.81, p=0.046) and validated (HR 1.43, 95%CI, 1.052-1.95, p=0.022) cohorts. Due to the small number of these studies meta-analysis was not carried out.

## Discussion

The results of the present systematic review and meta-analysis show clearly that the systemic inflammatory response, as evidenced by a number of markers at clinical thresholds, have independent prognostic value, across tumour types and geographical locations, in patients with advanced cancer. In particular, C-reactive protein, albumin and neutrophil count and the scores derived from them (GPS and NLR) have been consistently validated worldwide. There was considerable variation in the thresholds reported to have prognostic value when CRP, albumin and neutrophil counts were examined. There was less variation in the thresholds reported for NLR and still less for the GPS. The majority of studies were retrospective and therefore further prospective studies are warranted. In particular, there is a need to determine their clinical utility in the context of randomised clinical trials and thereby inform the appropriate treatment selection for patients with advanced cancer.

In the present review, the majority of studies reported overall survival as the end-point. However, for some markers of the systemic inflammatory response such as C-reactive protein and GPS there were also multiple studies using cancer specific survival as an end-point. It was of interest therefore that, on meta-analysis, the degree of heterogeneity appeared to be greater for overall survival as an endpoint compared with cancer specific survival (C-reactive protein 90% vs. 66% and GPS 56% vs. 4% respectively). This observation may be explained by previous observations that markers of the systemic inflammatory response have a stronger relationship with the cancer survival compared with the overall survival<sup>22,23</sup>. Therefore, it would appear that the optimal prognostic utility of markers of the systemic inflammatory response such as C-reactive protein and the GPS is in the prediction of cancer specific survival.

With reference to overall survival as an end-point, heterogeneity was greater in studies with a variety of thresholds compared to those with a standard threshold (e.g. C-reactive protein 90% (all) vs. 35% (>10mg/l), albumin 84% (all) vs. 79% (<35g/l) and NLR 79% (all) vs. 57% ( $\geq 5$ )).

respectively). In studies with these specific thresholds (e.g. in C-reactive protein threshold >10mg/l), compared with all tumour types, heterogeneity was less in specific tumour types (e.g. lung cancer heterogeneity was lower, 0% vs. 35% for all). Therefore, it would appear that the threshold used and the specific cancer studied influence the consistency of the association between markers of the systemic inflammatory response and overall survival in patients with advanced cancer. This has implications for the routine clinical application of markers such as C-reactive protein and NLR where a number of different thresholds have been reported in the literature. However, the GPS/mGPS have internationally recognised thresholds and are the preferred measure of the systemic inflammatory response amongst those investigators active in the field<sup>24</sup> and therefore are likely to have reproducible clinical utility in the context of randomized trials in patients with advanced cancer.

In the present review it was of interest that, across different markers of the systemic inflammatory response, when comparing using the same threshold and tumour type, the geographical prevalence of an elevated systemic inflammatory response varied. In particular, there was a trend towards a greater proportion of patients who had elevated markers in Western countries compared with Eastern Asian countries. Given the objective nature of these measurements there may be genetic or environmental causes of such a consistent difference. Indeed, there are well known ethnic differences in the normal range of neutrophils and lymphocytes<sup>25-27</sup>. Azab and co-workers recently reported that in a review of >9,000 patients, there were ethnic differences in NLR ratios in the United States<sup>27</sup>. Overall, the mean NLR was 2.15, whereas black Americans had a mean NLR of 1.76, Hispanic Americans had a mean NLR of 2.08 and white Americans had a mean NLR of 2.24<sup>27</sup>. Also, within ethnicities, patients who had diabetes, cardiovascular disease, a high BMI and were smokers had a significantly higher NLR<sup>27</sup>. Given that the most common thresholds used for NLR were >5 and >3 it is likely that a combination of genetic and environmental factors are responsible for such consistent East/West differences. To date, similar data for the GPS/mGPS has not appeared in the literature. Therefore, differences in the magnitude of systemic

inflammatory responses may explain, in part, the East/West split often observed in overall survival independent of tumour stage alone. Irrespective, the present results point to the value of not only staging the tumour but also the host systemic inflammatory response<sup>28</sup> in patients with advanced disease.

The systemic inflammatory response in patients with advanced cancer can be thought of as a result of a chronic inflammatory cascade. From the initial innate immune activation, as a result of the invasive tumour, due to the interaction of neutrophils and platelets at the site of tissue injury<sup>29</sup>, to the chronic wounding of tissues around the body in metastatic disease. This chronic activation of inflammatory processes results in profound changes at the genomic, intracellular, cellular and systemic levels in the patient with cancer<sup>9</sup>. In particular, at the systemic level, markers of a systemic inflammatory response are associated with a progressive nutritional and functional decline<sup>8</sup> and a profound deterioration in quality of life<sup>16</sup>.

A key pathway connecting the genomic, intracellular, cellular and systemic levels is the IL-6/JAK/STAT pathway<sup>30</sup>. Indeed, it is now increasingly recognised that genomic changes result in the chronic activation of the JAK/STAT pathway in the tumour and its microenvironment resulting in unregulated IL-6 production that produces an unregulated inflammatory cascade at cellular and systemic levels (increased C-reactive protein, neutrophil and platelet counts and decreased albumin). At the cellular and systemic level, IL-6 would appear to be the ideal marker of chronic systemic inflammation activation. Indeed, IL-6 in the circulation reflects the magnitude of tissue injury following surgery.<sup>31</sup> However, the strong correlation of IL-6 and C-reactive protein, the relative expense of IL-6 measurement has resulted in IL-6 not being routinely measured despite its central position in the systemic inflammatory cascade. Indeed, in the use of anti-IL-6 treatments, inhibition of the production and the fall in circulating concentrations of C-reactive protein is often used as a surrogate for IL-6 activity. Finally, that IL-6 is produced in most tissues including the tumour means that compared with C-reactive protein and albumin (produced in the liver only) and



The development of immune-oncology medications such as ipilimumab provides a potential means to target the activated inflammatory cascades to treat patients<sup>44:45</sup>. Indeed in a recent study in pancreatic cancer ruxolitinib, a strong downregulator of the inflammatory JAK/STAT pathway, was shown to increase median survival from 1.8 to 2.7 months in patients with high CRP readings<sup>46</sup>. This suggests a possible innovative means to treat patients with advanced cancers<sup>46</sup>.

The present systematic review and meta-analysis has a number of limitations. Intrinsic to the process and the high proportion of retrospective studies is the potential for publication bias. However, the volume of studies examined in the present review would mitigate, in part, against such publication bias. In the meta-analysis there was considerable heterogeneity that could be accounted for in part by differing thresholds and tumour type. It may be that as there is greater threshold standardisation in prospective studies the degree of heterogeneity will be reduced in subsequent meta-analysis of prospective studies.

In summary, the present systematic review and meta-analysis shows clearly that the systemic inflammatory response, as evidenced by a number of markers, has independent prognostic value in patients with advanced cancer. Of these markers, the GPS and NLR have been consistently validated worldwide. Therefore, it can be concluded that the systemic inflammatory response is an important predictor of outcome and is likely to inform treatment decisions in patients with advanced cancer. Further prospective studies are warranted.

The development of immune-oncology medications such as ipilimumab provides a potential means to target the activated inflammatory cascades to treat patients<sup>44;45</sup>. Indeed in a recent study in pancreatic cancer ruxolitinib, a strong downregulator of the inflammatory JAK/STAT pathway, was shown to increase median survival from 1.8 to 2.7 months in patients with high CRP readings<sup>46</sup>. This suggests a possible innovative means to treat patients with advanced cancers<sup>46</sup>.

The present systematic review and meta-analysis has a number of limitations. Intrinsic to the process and the high proportion of retrospective studies is the potential for publication bias. However, the volume of studies examined in the present review would mitigate, in part, against such publication bias. In the meta-analysis there was considerable heterogeneity that could be accounted for in part by differing thresholds and tumour type. It may be that as there is greater threshold standardisation in prospective studies the degree of heterogeneity will be reduced in subsequent meta-analysis of prospective studies.

In summary, the present systematic review and meta-analysis shows clearly that the systemic inflammatory response, as evidenced by a number of markers, has independent prognostic value in patients with advanced cancer. Of these markers, the GPS and NLR have been consistently validated worldwide. Therefore, it can be concluded that the systemic inflammatory response is an important predictor of outcome and is likely to inform treatment decisions in patients with advanced cancer. Further prospective studies are warranted.

## Reference List

- (1) Garraway LA, Verweij J, Ballman KV. Precision oncology: an overview. *J Clin Oncol* 2013; **31**(15):1803-1805.
- (2) Prigerson HG, Bao Y, Shah MA, Paulk ME, LeBlanc TW, Schneider BJ et al. Chemotherapy Use, Performance Status, and Quality of Life at the End of Life. *JAMA Oncol* 2015; **1**(6):778-784.
- (3) Garrido MM, Prigerson HG, Bao Y, Maciejewski PK. Chemotherapy Use in the Months Before Death and Estimated Costs of Care in the Last Week of Life. *J Pain Symptom Manage* 2016; **51**(5):875-881.
- (4) Mayor S. UK audit shows need for greater caution with chemotherapy in very sick patients. *BMJ* 2008; **337**:a2498.
- (5) Temel JS, Greer JA, Muzikansky A, Gallagher ER, Admane S, Jackson VA et al. Early palliative care for patients with metastatic non-small-cell lung cancer. *N Engl J Med* 2010; **363**(8):733-742.
- (6) Martin L, Birdsell L, MacDonald N, Reiman T, Clandinin MT, McCargar LJ et al. Cancer cachexia in the age of obesity: skeletal muscle depletion is a powerful prognostic factor, independent of body mass index. *J Clin Oncol* 2013; **31**(12):1539-1547.
- (7) Simmons CP, McMillan DC, McWilliams K, Sande TA, Fearon KC, Tuck S et al. Prognostic Tools in Patients with Advanced Cancer: A Systematic Review. *J Pain Symptom Manage* 2017.
- (8) McMillan DC. Systemic inflammation, nutritional status and survival in patients with cancer. *Curr Opin Clin Nutr Metab Care* 2009; **12**(3):223-226.
- (9) Diakos CI, Charles KA, McMillan DC, Clarke SJ. Cancer-related inflammation and treatment effectiveness. *Lancet Oncol* 2014; **15**(11):e493-e503.
- (10) Johns N, Hatakeyama S, Stephens NA, Degen M, Degen S, Frieauff W et al. Clinical classification of cancer cachexia: phenotypic correlates in human skeletal muscle. *PLoS One* 2014; **9**(1):e83618.
- (11) Kerem M, Ferahkose Z, Yilmaz UT, Pasaoglu H, Ofluoglu E, Bedirli A et al. Adipokines and ghrelin in gastric cancer cachexia. *World J Gastroenterol* 2008; **14**(23):3633-3641.
- (12) Guthrie GJ, Charles KA, Roxburgh CS, Horgan PG, McMillan DC, Clarke SJ. The systemic inflammation-based neutrophil-lymphocyte ratio: experience in patients with cancer. *Crit Rev Oncol Hematol* 2013; **88**(1):218-230.
- (13) McMillan DC. An inflammation-based prognostic score and its role in the nutrition-based management of patients with cancer. *Proc Nutr Soc* 2008; **67**(3):257-262.
- (14) McMillan DC. The systemic inflammation-based Glasgow Prognostic Score: a decade of experience in patients with cancer. *Cancer Treat Rev* 2013; **39**(5):534-540.

- (15) Laird BJ, Kaasa S, McMillan DC, Fallon MT, Hjermstad MJ, Fayers P et al. Prognostic factors in patients with advanced cancer: a comparison of clinicopathological factors and the development of an inflammation-based prognostic system. *Clin Cancer Res* 2013; **19**(19):5456-5464.
- (16) Laird BJ, Fallon M, Hjermstad MJ, Tuck S, Kaasa S, Klepstad P et al. Quality of Life in Patients With Advanced Cancer: Differential Association With Performance Status and Systemic Inflammatory Response. *J Clin Oncol* 2016; **34**(23):2769-2775.
- (17) Zhou T, Zhan J, Hong S, Hu Z, Fang W, Qin T et al. Ratio of C-Reactive Protein/Albumin is An Inflammatory Prognostic Score for Predicting Overall Survival of Patients with Small-cell Lung Cancer. *Sci Rep* 2015; **5**:10481.
- (18) Yamashita S, Kohjimoto Y, Iguchi T, Koike H, Kusumoto H, Iba A et al. Prognostic factors and risk stratification in patients with castration-resistant prostate cancer receiving docetaxel-based chemotherapy. *BMC Urol* 2016; **16**:13.
- (19) Shibutani M, Maeda K, Nagahara H, Ohtani H, Iseki Y, Ikeya T et al. The pretreatment albumin to globulin ratio predicts chemotherapeutic outcomes in patients with unresectable metastatic colorectal cancer. *BMC Cancer* 2015; **15**:347.
- (20) Yao Y, Zhao M, Yuan D, Gu X, Liu H, Song Y. Elevated pretreatment serum globulin albumin ratio predicts poor prognosis for advanced non-small cell lung cancer patients. *J Thorac Dis* 2014; **6**(9):1261-1270.
- (21) Chan AW, Chan SL, Mo FK, Wong GL, Wong VW, Cheung YS et al. Albumin-to-alkaline phosphatase ratio: a novel prognostic index for hepatocellular carcinoma. *Dis Markers* 2015; **2015**:564057.
- (22) Marsik C, Kazemi-Shirazi L, Schickbauer T, Winkler S, Joukhadar C, Wagner OF et al. C-reactive protein and all-cause mortality in a large hospital-based cohort. *Clin Chem* 2008; **54**(2):343-349.
- (23) Proctor MJ, McMillan DC, Horgan PG, Fletcher CD, Talwar D, Morrison DS. Systemic inflammation predicts all-cause mortality: a glasgow inflammation outcome study. *PLoS One* 2015; **10**(3):e0116206.
- (24) Watt DG, Roxburgh CS, White M, Chan JZ, Horgan PG, McMillan DC. A Survey of Attitudes towards the Clinical Application of Systemic Inflammation Based Prognostic Scores in Cancer. *Mediators Inflamm* 2015; **2015**:842070.
- (25) Bain B, Seed M, Godsland I. Normal values for peripheral blood white cell counts in women of four different ethnic origins. *J Clin Pathol* 1984; **37**(2):188-193.
- (26) Bain BJ. Ethnic and sex differences in the total and differential white cell count and platelet count. *J Clin Pathol* 1996; **49**(8):664-666.
- (27) Azab B, Camacho-Rivera M, Taioli E. Average values and racial differences of neutrophil lymphocyte ratio among a nationally representative sample of United States subjects. *PLoS One* 2014; **9**(11):e112361.

- (28) McMillan DC. Cancer and systemic inflammation: stage the tumour and stage the host. *Br J Cancer* 2013; **109**(3):529.
- (29) Sreeramkumar V, Adrover JM, Ballesteros I, Cuartero MI, Rossaint J, Bilbao I et al. Neutrophils scan for activated platelets to initiate inflammation. *Science* 2014; **346**(6214):1234-1238.
- (30) Roxburgh CS, McMillan DC. Therapeutics targeting innate immune/inflammatory responses through the interleukin-6/JAK/STAT signal transduction pathway in patients with cancer. *Transl Res* 2016; **167**(1):61-66.
- (31) Watt DG, Horgan PG, McMillan DC. Routine clinical markers of the magnitude of the systemic inflammatory response after elective operation: a systematic review. *Surgery* 2015; **157**(2):362-380.
- (32) Crozier JE, McKee RF, McArdle CS, Angerson WJ, Anderson JH, Horgan PG et al. The presence of a systemic inflammatory response predicts poorer survival in patients receiving adjuvant 5-FU chemotherapy following potentially curative resection for colorectal cancer. *Br J Cancer* 2006; **94**(12):1833-1836.
- (33) Carruthers R, Tho LM, Brown J, Kakumanu S, McCartney E, McDonald AC. Systemic inflammatory response is a predictor of outcome in patients undergoing preoperative chemoradiation for locally advanced rectal cancer. *Colorectal Dis* 2012; **14**(10):e701-e707.
- (34) Dreyer SB, Powell AG, McSorley ST, Waterston A, Going JJ, Edwards J et al. The Pretreatment Systemic Inflammatory Response is an Important Determinant of Poor Pathologic Response for Patients Undergoing Neoadjuvant Therapy for Rectal Cancer. *Ann Surg Oncol* 2017; **24**(5):1295-1303.
- (35) McSorley ST, Dolan RD, Roxburgh CS, McMillan DC, Horgan PG. How and why systemic inflammation worsens quality of life in patients with advanced cancer. 14-2-2017.

Ref Type: Online Source

- (36) Lee BN, Dantzer R, Langley KE, Bennett GJ, Dougherty PM, Dunn AJ et al. A cytokine-based neuroimmunologic mechanism of cancer-related symptoms. *Neuroimmunomodulation* 2004; **11**(5):279-292.
- (37) De PM, Lewis CE. Macrophage regulation of tumor responses to anticancer therapies. *Cancer Cell* 2013; **23**(3):277-286.
- (38) Wood LJ, Nail LM, Gilster A, Winters KA, Elsea CR. Cancer chemotherapy-related symptoms: evidence to suggest a role for proinflammatory cytokines. *Oncol Nurs Forum* 2006; **33**(3):535-542.
- (39) Wood LJ, Nail LM, Perrin NA, Elsea CR, Fischer A, Druker BJ. The cancer chemotherapy drug etoposide (VP-16) induces proinflammatory cytokine production and sickness behavior-like symptoms in a mouse model of cancer chemotherapy-related symptoms. *Biol Res Nurs* 2006; **8**(2):157-169.

- (40) Tsavaris N, Kosmas C, Vadiaka M, Kanelopoulos P, Boulamatsis D. Immune changes in patients with advanced breast cancer undergoing chemotherapy with taxanes. *Br J Cancer* 2002; **87**(1):21-27.
- (41) Plate JM, Plate AE, Shott S, Bograd S, Harris JE. Effect of gemcitabine on immune cells in subjects with adenocarcinoma of the pancreas. *Cancer Immunol Immunother* 2005; **54**(9):915-925.
- (42) Pusztai L, Mendoza TR, Reuben JM, Martinez MM, Willey JS, Lara J et al. Changes in plasma levels of inflammatory cytokines in response to paclitaxel chemotherapy. *Cytokine* 2004; **25**(3):94-102.
- (43) Wang XS, Shi Q, Williams LA, Mao L, Cleeland CS, Komaki RR et al. Inflammatory cytokines are associated with the development of symptom burden in patients with NSCLC undergoing concurrent chemoradiation therapy. *Brain Behav Immun* 2010; **24**(6):968-974.
- (44) Hoos A. Development of immuno-oncology drugs - from CTLA4 to PD1 to the next generations. *Nat Rev Drug Discov* 2016; **15**(4):235-247.
- (45) Hoos A, Ibrahim R, Korman A, Abdallah K, Berman D, Shahabi V et al. Development of ipilimumab: contribution to a new paradigm for cancer immunotherapy. *Semin Oncol* 2010; **37**(5):533-546.
- (46) Ruxolitinib Benefits Some with Pancreatic Cancer. *Cancer Discov* 2015; **5**(12):1231.
- (47) Ueno H, Okada S, Okusaka T, Ikeda M. Prognostic factors in patients with metastatic pancreatic adenocarcinoma receiving systemic chemotherapy. *Oncology* 2000; **59**(4):296-301.
- (48) McMillan DC, Elahi MM, Sattar N, Angerson WJ, Johnstone J, McArdle CS. Measurement of the systemic inflammatory response predicts cancer-specific and non-cancer survival in patients with cancer. *Nutr Cancer* 2001; **41**(1-2):64-69.
- (49) Scott HR, McMillan DC, Forrest LM, Brown DJ, McArdle CS, Milroy R. The systemic inflammatory response, weight loss, performance status and survival in patients with inoperable non-small cell lung cancer. *Br J Cancer* 2002; **87**(3):264-267.
- (50) Bromwich E, McMillan DC, Lamb GW, Vasey PA, Aitchison M. The systemic inflammatory response, performance status and survival in patients undergoing alpha-interferon treatment for advanced renal cancer. *Br J Cancer* 2004; **91**(7):1236-1238.
- (51) Elahi MM, McMillan DC, McArdle CS, Angerson WJ, Soukop M, Johnstone J et al. The systemic inflammatory response predicts overall and cancer specific survival in patients with malignant lymphoma. *Med Sci Monit* 2005; **11**(2):CR75-CR78.
- (52) Casamassima A, Picciariello M, Quaranta M, Berardino R, Ranieri C, Paradiso A et al. C-reactive protein: a biomarker of survival in patients with metastatic renal cell carcinoma treated with subcutaneous interleukin-2 based immunotherapy. *J Urol* 2005; **173**(1):52-55.

- (53) McArdle PA, Mir K, Almushatat AS, Wallace AM, Underwood MA, McMillan DC. Systemic inflammatory response, prostate-specific antigen and survival in patients with metastatic prostate cancer. *Urol Int* 2006; **77**(2):127-129.
- (54) Sawaki A, Kanemitsu Y, Mizuno N, Takahashi K, Nakamura T, Ioka T et al. Practical prognostic index for patients with metastatic pancreatic cancer treated with gemcitabine. *J Gastroenterol Hepatol* 2008; **23**(8 Pt 1):1292-1297.
- (55) Al Murri AM, Bartlett JM, Canney PA, Doughty JC, Wilson C, McMillan DC. Evaluation of an inflammation-based prognostic score (GPS) in patients with metastatic breast cancer. *Br J Cancer* 2006; **94**(2):227-230.
- (56) Nakachi K, Furuse J, Ishii H, Suzuki E, Yoshino M. Prognostic factors in patients with gemcitabine-refractory pancreatic cancer. *Jpn J Clin Oncol* 2007; **37**(2):114-120.
- (57) Ramsey S, Lamb GW, Aitchison M, Graham J, McMillan DC. Evaluation of an inflammation-based prognostic score in patients with metastatic renal cancer. *Cancer* 2007; **109**(2):205-212.
- (58) Tanaka T, Ikeda M, Okusaka T, Ueno H, Morizane C, Hagihara A et al. Prognostic factors in Japanese patients with advanced pancreatic cancer treated with single-agent gemcitabine as first-line therapy. *Jpn J Clin Oncol* 2008; **38**(11):755-761.
- (59) Beer TM, Lalani AS, Lee S, Mori M, Eilers KM, Curd JG et al. C-reactive protein as a prognostic marker for men with androgen-independent prostate cancer: results from the ASCENT trial. *Cancer* 2008; **112**(11):2377-2383.
- (60) Papadoniou N, Kosmas C, Gennatas K, Polyzos A, Mouratidou D, Skopelitis E et al. Prognostic factors in patients with locally advanced (unresectable) or metastatic pancreatic adenocarcinoma: a retrospective analysis. *Anticancer Res* 2008; **28**(1B):543-549.
- (61) Yoshida S, Saito K, Koga F, Yokoyama M, Kageyama Y, Masuda H et al. C-reactive protein level predicts prognosis in patients with muscle-invasive bladder cancer treated with chemoradiotherapy. *BJU Int* 2008; **101**(8):978-981.
- (62) Koch A, Fohlin H, Sorenson S. Prognostic significance of C-reactive protein and smoking in patients with advanced non-small cell lung cancer treated with first-line palliative chemotherapy. *J Thorac Oncol* 2009; **4**(3):326-332.
- (63) Hashimoto K, Ueno H, Ikeda M, Kojima Y, Hagihara A, Kondo S et al. Do recurrent and metastatic pancreatic cancer patients have the same outcomes with gemcitabine treatment? *Oncology* 2009; **77**(3-4):217-223.
- (64) Zacharakis M, Xynos ID, Lazaris A, Smaro T, Kosmas C, Dokou A et al. Predictors of survival in stage IV metastatic colorectal cancer. *Anticancer Res* 2010; **30**(2):653-660.
- (65) Iwasa S, Nakajima TE, Nakamura K, Takashima A, Kato K, Hamaguchi T et al. Systemic chemotherapy for peritoneal disseminated gastric cancer with inadequate oral intake: a retrospective study. *Int J Clin Oncol* 2011; **16**(1):57-62.

- (66) Falkensammer CE, Thurnher M, Leonhartsberger N, Ramoner R. C-reactive protein is a strong predictor for anaemia in renal cell carcinoma: role of IL-6 in overall survival. *BJU Int* 2011; **107**(12):1893-1898.
- (67) Masago K, Fujita S, Togashi Y, Kim YH, Hatachi Y, Fukuhara A et al. Clinical significance of pretreatment C-reactive protein in patients with advanced nonsquamous, non-small cell lung cancer who received gefitinib. *Oncology* 2010; **79**(5-6):355-362.
- (68) Shimoda M, Katoh M, Kita J, Sawada T, Kubota K. The Glasgow Prognostic Score is a good predictor of treatment outcome in patients with unresectable pancreatic cancer. *Chemotherapy* 2010; **56**(6):501-506.
- (69) Shinohara N, Abe T, Mochizuki T, Kashiwagi A, Kanagawa K, Maruyama S et al. Is Memorial Sloan-Kettering Cancer Center risk classification appropriate for Japanese patients with metastatic renal cell carcinoma in the cytokine era? *Urol Oncol* 2013; **31**(7):1276-1282.
- (70) Yi JH, Lee J, Park SH, Lee KT, Lee JK, Lee KH et al. A prognostic model to predict clinical outcomes with first-line gemcitabine-based chemotherapy in advanced pancreatic cancer. *Oncology* 2011; **80**(3-4):175-180.
- (71) Kume H, Kakutani S, Yamada Y, Shinohara M, Tominaga T, Suzuki M et al. Prognostic factors for renal cell carcinoma with bone metastasis: who are the long-term survivors? *J Urol* 2011; **185**(5):1611-1614.
- (72) Lee JS, Kwon OY, Choi HS, Hong HP, Ko YG. Serum C-reactive protein level is a predictive factor for 14-day mortality of patients with advanced cancer who present to the emergency department with acute symptoms. *Acad Emerg Med* 2011; **18**(4):440-442.
- (73) Bystrom P, Berglund A, Nygren P, Wernroth L, Johansson B, Larsson A et al. Evaluation of predictive markers for patients with advanced colorectal cancer. *Acta Oncol* 2012; **51**(7):849-859.
- (74) Ishioka J, Saito K, Sakura M, Yokoyama M, Matsuoka Y, Numao N et al. Development of a nomogram incorporating serum C-reactive protein level to predict overall survival of patients with advanced urothelial carcinoma and its evaluation by decision curve analysis. *Br J Cancer* 2012; **107**(7):1031-1036.
- (75) Prins RC, Rademacher BL, Mongoue-Tchokote S, Alumkal JJ, Graff JN, Eilers KM et al. C-reactive protein as an adverse prognostic marker for men with castration-resistant prostate cancer (CRPC): confirmatory results. *Urol Oncol* 2012; **30**(1):33-37.
- (76) Zeng YC, Xue M, Chi F, Xu ZG, Fan GL, Wu R et al. C-reactive protein level predicts prognosis in patients with locoregionally advanced laryngeal carcinoma treated with chemoradiotherapy. *Tumour Biol* 2012; **33**(3):891-895.
- (77) Pond GR, Armstrong AJ, Wood BA, Leopold L, Galsky MD, Sonpavde G. Ability of C-reactive protein to complement multiple prognostic classifiers in men with metastatic castration resistant prostate cancer receiving docetaxel-based chemotherapy. *BJU Int* 2012; **110**(11 Pt B):E461-E468.



- (78) Kinoshita A, Onoda H, Takano K, Imai N, Saeki C, Fushiya N et al. Pretreatment serum C-reactive protein level predicts poor prognosis in patients with hepatocellular carcinoma. *Med Oncol* 2012; **29**(4):2800-2808.
- (79) Morizane S, Iwamoto H, Yao A, Isoyama T, Sejima T, Takenaka A. Serum C-reactive protein level is a significant prognostic indicator in patients with advanced urothelial cancer treated with gemcitabine-cisplatin or carboplatin - preliminary results. *Cent European J Urol* 2012; **65**(2):62-66.
- (80) Haas M, Heinemann V, Kullmann F, Laubender RP, Klose C, Bruns CJ et al. Prognostic value of CA 19-9, CEA, CRP, LDH and bilirubin levels in locally advanced and metastatic pancreatic cancer: results from a multicenter, pooled analysis of patients receiving palliative chemotherapy. *J Cancer Res Clin Oncol* 2013; **139**(4):681-689.
- (81) Xia WX, Zhang HB, Shi JL, Lu X, Wang L, Ye YF et al. A prognostic model predicts the risk of distant metastasis and death for patients with nasopharyngeal carcinoma based on pre-treatment serum C-reactive protein and N-classification. *Eur J Cancer* 2013; **49**(9):2152-2160.
- (82) Yasuda Y, Saito K, Yuasa T, Kitsukawa S, Urakami S, Yamamoto S et al. Prognostic impact of pretreatment C-reactive protein for patients with metastatic renal cell carcinoma treated with tyrosine kinase inhibitors. *Int J Clin Oncol* 2013; **18**(5):884-889.
- (83) Shirakawa T, Kato K, Nagashima K, Nishikawa A, Sawada R, Takahashi N et al. A retrospective study of docetaxel or paclitaxel in patients with advanced or recurrent esophageal squamous cell carcinoma who previously received fluoropyrimidine- and platinum-based chemotherapy. *Cancer Chemother Pharmacol* 2014; **74**(6):1207-1215.
- (84) Teishima J, Kobatake K, Hayashi T, Seno Y, Ikeda K, Nagamatsu H et al. Prognostic significance of C-reactive protein in patients with intermediate-risk metastatic renal cell carcinoma treated with molecular targeted therapy. *Oncol Lett* 2014; **8**(2):881-885.
- (85) Deberne M, Ropert S, Billemont B, Daniel C, Chapron J, Goldwasser F. Inaugural bone metastases in non-small cell lung cancer: a specific prognostic entity? *BMC Cancer* 2014; **14**:416.
- (86) Beuselinck B, Vano YA, Oudard S, Wolter P, De SR, Depoorter L et al. Prognostic impact of baseline serum C-reactive protein in patients with metastatic renal cell carcinoma (RCC) treated with sunitinib. *BJU Int* 2014; **114**(1):81-89.
- (87) Xue P, Kanai M, Mori Y, Nishimura T, Uza N, Kodama Y et al. Comparative outcomes between initially unresectable and recurrent cases of advanced pancreatic cancer following palliative chemotherapy. *Pancreas* 2014; **43**(3):411-416.
- (88) Formica V, Luccchetti J, Cunningham D, Smyth EC, Ferroni P, Nardecchia A et al. Systemic inflammation, as measured by the neutrophil/lymphocyte ratio, may have differential prognostic impact before and during treatment with fluorouracil, irinotecan and bevacizumab in metastatic colorectal cancer patients. *Med Oncol* 2014; **31**(9):166.
- (89) Kim YJ, Kim SJ, Lee JK, Choi WS, Park JH, Kim HJ et al. Prediction of survival in terminally ill cancer patients at the time of terminal cancer diagnosis. *J Cancer Res Clin Oncol* 2014; **140**(9):1567-1574.

- (90) Ni XF, Wu P, Wu CP, Ji M, Wu J, Gu XF et al. Elevated serum C-reactive protein, carcinoembryonic antigen and N2 disease are poor prognostic indicators in non-small cell lung cancer. *Asia Pac J Clin Oncol* 2015; **11**(4):e22-e30.
- (91) Fiala O, Pesek M, Finek J, Topolcan O, Racek J, Minarik M et al. High serum level of C-reactive protein is associated with worse outcome of patients with advanced-stage NSCLC treated with erlotinib. *Tumour Biol* 2015; **36**(12):9215-9222.
- (92) Adams HJ, de Klerk JM, Fijnheer R, Heggelman BG, Dubois SV, Nievelstein RA et al. Prognostic Value of Anemia and C-Reactive Protein Levels in Diffuse Large B-Cell Lymphoma. *Clin Lymphoma Myeloma Leuk* 2015; **15**(11):671-679.
- (93) Ito M, Saito K, Yasuda Y, Sukegawa G, Kubo Y, Numao N et al. Prognostic impact of C-reactive protein for determining overall survival of patients with castration-resistant prostate cancer treated with docetaxel. *Urology* 2011; **78**(5):1131-1135.
- (94) Li X, Tian F, Wang F, Li Y. Serum C-reactive protein and overall survival of patients with osteosarcoma. *Tumour Biol* 2015; **36**(7):5663-5666.
- (95) Tang LQ, Hu DP, Chen QY, Zhang L, Lai XP, He Y et al. Elevated high-sensitivity C-reactive protein levels predict decreased survival for nasopharyngeal carcinoma patients in the intensity-modulated radiotherapy era. *PLoS One* 2015; **10**(4):e0122965.
- (96) Thurner EM, Krenn-Pilko S, Langsenlehner U, Stojakovic T, Pichler M, Gerger A et al. The elevated C-reactive protein level is associated with poor prognosis in prostate cancer patients treated with radiotherapy. *Eur J Cancer* 2015; **51**(5):610-619.
- (97) Zeng YC, Wu R, Xiao YP, Chi F, Xue M, Zhang ZY et al. Serum C-reactive protein predicts poor prognosis in patients with locoregionally advanced nasopharyngeal carcinoma treated with chemoradiotherapy. *Curr Oncol* 2015; **22**(1):20-24.
- (98) Xu L, Zhao Q, Huang S, Li S, Wang J, Li Q. Serum C-reactive protein acted as a prognostic biomarker for overall survival in metastatic prostate cancer patients. *Tumour Biol* 2015; **36**(2):669-673.
- (99) Go SI, Kim RB, Song HN, Kang MH, Lee US, Choi HJ et al. Prognostic significance of the absolute monocyte counts in lung cancer patients with venous thromboembolism. *Tumour Biol* 2015; **36**(10):7631-7639.
- (100) Martin HL, Ohara K, Kiberu A, Van HT, Davidson A, Khattak MA. Prognostic value of systemic inflammation-based markers in advanced pancreatic cancer. *Intern Med J* 2014; **44**(7):676-682.
- (101) Mitsunaga S, Ikeda M, Shimizu S, Ohno I, Takahashi H, Okuyama H et al. C-Reactive Protein Level Is an Indicator of the Aggressiveness of Advanced Pancreatic Cancer. *Pancreas* 2016; **45**(1):110-116.
- (102) Kim HW, Lee JC, Paik KH, Lee YS, Hwang JH, Kim J. Initial Metastatic Site as a Prognostic Factor in Patients With Stage IV Pancreatic Ductal Adenocarcinoma. *Medicine (Baltimore)* 2015; **94**(25):e1012.

- (103) Yao A, Sejima T, Iwamoto H, Masago T, Morizane S, Honda M et al. High neutrophil-to-lymphocyte ratio predicts poor clinical outcome in patients with castration-resistant prostate cancer treated with docetaxel chemotherapy. *Int J Urol* 2015; **22**(9):827-833.
- (104) Wu G, Yao Y, Bai C, Zeng J, Shi D, Gu X et al. Combination of platelet to lymphocyte ratio and neutrophil to lymphocyte ratio is a useful prognostic factor in advanced non-small cell lung cancer patients. *Thorac Cancer* 2015; **6**(3):275-287.
- (105) Middleton G, Greenhalf W, Costello E, Shaw V, Cox T, Ghaneh P et al. Immunobiological effects of gemcitabine and capecitabine combination chemotherapy in advanced pancreatic ductal adenocarcinoma. *Br J Cancer* 2016; **114**(5):510-518.
- (106) Casadei GA, Carloni S, Scarpi E, Maltoni P, Dorizzi RM, Passardi A et al. Prognostic role of serum concentrations of high-sensitivity C-reactive protein in patients with metastatic colorectal cancer: results from the ITACa trial. *Oncotarget* 2016; **7**(9):10193-10202.
- (107) Sheng J, Yang YP, Ma YX, Qin T, Hu ZH, Hong SD et al. Low Prognostic Nutritional Index Correlates with Worse Survival in Patients with Advanced NSCLC following EGFR-TKIs. *PLoS One* 2016; **11**(1):e0147226.
- (108) Kou T, Kanai M, Yamamoto M, Xue P, Mori Y, Kudo Y et al. Prognostic model for survival based on readily available pretreatment factors in patients with advanced pancreatic cancer receiving palliative chemotherapy. *Int J Clin Oncol* 2016; **21**(1):118-125.
- (109) Ahn HK, Hwang IC, Lee JS, Sym SJ, Cho EK, Shin DB. Neutrophil-Lymphocyte Ratio Predicts Survival in Terminal Cancer Patients. *J Palliat Med* 2016; **19**(4):437-441.
- (110) Axedorph U, Sjoberg J, Grimfors G, Landgren O, Porwit-Macdonald A, Bjorkholm M. Biological markers may add to prediction of outcome achieved by the International Prognostic Score in Hodgkin's disease. *Ann Oncol* 2000; **11**(11):1405-1411.
- (111) Vigano A, Bruera E, Jhangri GS, Newman SC, Fields AL, Suarez-Almazor ME. Clinical survival predictors in patients with advanced cancer. *Arch Intern Med* 2000; **160**(6):861-868.
- (112) Marechal R, Demols A, Gay F, De M, V, Arvanitaki M, Hendlisz A et al. Prognostic factors and prognostic index for chemonaive and gemcitabine-refractory patients with advanced pancreatic cancer. *Oncology* 2007; **73**(1-2):41-51.
- (113) Lam PT, Leung MW, Tse CY. Identifying prognostic factors for survival in advanced cancer patients: a prospective study. *Hong Kong Med J* 2007; **13**(6):453-459.
- (114) Paralkar VR, Li T, Langer CJ. Population characteristics and prognostic factors in metastatic non-small-cell lung cancer: a Fox Chase Cancer Center retrospective. *Clin Lung Cancer* 2008; **9**(2):116-121.
- (115) Ngo L, Hee SW, Lim LC, Tao M, Quek R, Yap SP et al. Prognostic factors in patients with diffuse large B cell lymphoma: Before and after the introduction of rituximab. *Leuk Lymphoma* 2008; **49**(3):462-469.

- (116) Shim HJ, Yun JY, Hwang JE, Bae WK, Cho SH, Chung JJ. Prognostic factor analysis of third-line chemotherapy in patients with advanced gastric cancer. *Gastric Cancer* 2011; **14**(3):249-256.
- (117) Tredan O, Ray-Coquard I, Chvetzoff G, Rebattu P, Bajard A, Chabaud S et al. Validation of prognostic scores for survival in cancer patients beyond first-line therapy. *BMC Cancer* 2011; **11**:95.
- (118) Lim KH, Han SW, Oh DY, Im SA, Kim TY, Bang YJ. Outcome of infusional 5-fluorouracil, doxorubicin, and mitomycin-C (iFAM) chemotherapy and analysis of prognostic factors in patients with refractory advanced biliary tract cancer. *Oncology* 2012; **83**(2):57-66.
- (119) Prakash G, Sharma A, Raina V, Kumar L, Sharma MC, Mohanti BK. B cell non-Hodgkin's lymphoma: experience from a tertiary care cancer center. *Ann Hematol* 2012; **91**(10):1603-1611.
- (120) Kang EJ, Choi YJ, Kim JS, Park KH, Oh SC, Seo JH et al. Prognostic Factors for the Selection of Patients Eligible for Second-Line Chemotherapy in Advanced Biliary Tract Cancer. *Chemotherapy* 2014; **60**(2):91-98.
- (121) Ulas A, Turkoz FP, Silay K, Tokluoglu S, Avci N, Oksuzoglu B et al. A laboratory prognostic index model for patients with advanced non-small cell lung cancer. *PLoS One* 2014; **9**(12):e114471.
- (122) Imedio ER, Beveridge RD, Urtasun JA, Campos GB, Estelles DL, Esparcia MF et al. Safety and efficacy of sorafenib in the treatment of advanced hepatocellular carcinoma: a single center experience. *Med Oncol* 2014; **31**(5):948.
- (123) Malik L, Parsons H, Mahalingam D, Ehler B, Goros M, Mejia A et al. Clinical outcomes and survival of advanced renal cancer patients in phase I clinical trials. *Clin Genitourin Cancer* 2014; **12**(5):359-365.
- (124) Tsai HJ, Hsieh MY, Tsai YC, Liu ZY, Hsieh HY, Lee CM et al. Liver function tests may be useful tools for advanced cancer patient care: a preliminary single-center result. *Kaohsiung J Med Sci* 2014; **30**(3):146-152.
- (125) Stenman M, Laurell A, Lindskog M. Prognostic significance of serum albumin in patients with metastatic renal cell carcinoma. *Med Oncol* 2014; **31**(3):841.
- (126) Koo DH, Ryu MH, Ryoo BY, Seo J, Lee MY, Chang HM et al. Improving trends in survival of patients who receive chemotherapy for metastatic or recurrent gastric cancer: 12 years of experience at a single institution. *Gastric Cancer* 2015; **18**(2):346-353.
- (127) Kao J, Gold KD, Zarrili G, Copel E, Silverman AJ, Ramsaran SS et al. Clinical Predictors of Survival for Patients with Stage IV Cancer Referred to Radiation Oncology. *PLoS One* 2015; **10**(4):e0124329.
- (128) Wild AT, Ye X, Ellsworth SG, Smith JA, Narang AK, Garg T et al. The Association Between Chemoradiation-related Lymphopenia and Clinical Outcomes in Patients With Locally Advanced Pancreatic Adenocarcinoma. *Am J Clin Oncol* 2015; **38**(3):259-265.

- (129) Helissey C, Berger F, Cottu P, Dieras V, Mignot L, Servois V et al. Circulating tumor cell thresholds and survival scores in advanced metastatic breast cancer: the observational step of the CirCe01 phase III trial. *Cancer Lett* 2015; **360**(2):213-218.
- (130) Narwani V, Gabriel J, Boyd K, Chevassut T. Absolute lymphocyte count at day 29 of treatment is a powerful predictor of outcome in multiple myeloma. *Clin Lymphoma Myeloma Leuk* 2015; **15**(4):222-226.
- (131) Moon H, Roh JL, Lee SW, Kim SB, Choi SH, Nam SY et al. Prognostic value of nutritional and hematologic markers in head and neck squamous cell carcinoma treated by chemoradiotherapy. *Radiother Oncol* 2016; **118**(2):330-334.
- (132) Uemura K, Miyoshi Y, Kawahara T, Yoneyama S, Hattori Y, Teranishi J et al. Prognostic value of a computer-aided diagnosis system involving bone scans among men treated with docetaxel for metastatic castration-resistant prostate cancer. *BMC Cancer* 2016; **16**:109.
- (133) Dorajoo SR, Tan WJ, Koo SX, Tan WS, Chew MH, Tang CL et al. A scoring model for predicting survival following primary tumour resection in stage IV colorectal cancer patients with unresectable metastasis. *Int J Colorectal Dis* 2016; **31**(2):235-245.
- (134) Choi Y, Oh DY, Park H, Kim TY, Lee KH, Han SW et al. More Accurate Prediction of Metastatic Pancreatic Cancer Patients' Survival with Prognostic Model Using Both Host Immunity and Tumor Metabolic Activity. *PLoS One* 2016; **11**(1):e0145692.
- (135) Mandrekar SJ, Schild SE, Hillman SL, Allen KL, Marks RS, Mailliard JA et al. A prognostic model for advanced stage nonsmall cell lung cancer. Pooled analysis of North Central Cancer Treatment Group trials. *Cancer* 2006; **107**(4):781-792.
- (136) Tibaldi C, Vasile E, Bernardini I, Orlandini C, Andreuccetti M, Falcone A. Baseline elevated leukocyte count in peripheral blood is associated with poor survival in patients with advanced non-small cell lung cancer: a prognostic model. *J Cancer Res Clin Oncol* 2008; **134**(10):1143-1149.
- (137) Partridge M, Fallon M, Bray C, McMillan D, Brown D, Laird B. Prognostication in advanced cancer: a study examining an inflammation-based score. *J Pain Symptom Manage* 2012; **44**(2):161-167.
- (138) Lee CK, Hudson M, Stockler M, Coates AS, Ackland S, Gebski V et al. A nomogram to predict survival time in women starting first-line chemotherapy for advanced breast cancer. *Breast Cancer Res Treat* 2011; **129**(2):467-476.
- (139) Kawashima A, Tsujimura A, Takayama H, Arai Y, Nin M, Tanigawa G et al. Impact of hyponatremia on survival of patients with metastatic renal cell carcinoma treated with molecular targeted therapy. *Int J Urol* 2012; **19**(12):1050-1057.
- (140) Luo XL, He W, Huang H, Chen QY, Liang Y, Mai HQ et al. Design of a prognostic score model for nasopharyngeal carcinoma. *Head Neck* 2015; **37**(5):624-629.
- (141) Iacovelli R, Farcomeni A, Sternberg CN, Carteni G, Milella M, Santoni M et al. Prognostic factors in patients receiving third line targeted therapy for metastatic renal cell carcinoma. *J Urol* 2015; **193**(6):1905-1910.

- (142) Ferrucci PF, Ascierto PA, Pigozzo J, Del VM, Maio M, Antonini Cappellini GC et al. Baseline neutrophils and derived neutrophil-to-lymphocyte ratio: prognostic relevance in metastatic melanoma patients receiving ipilimumab. *Ann Oncol* 2016; **27**(4):732-738.
- (143) Bille A, Krug LM, Woo KM, Rusch VW, Zauderer MG. Contemporary Analysis of Prognostic Factors in Patients with Unresectable Malignant Pleural Mesothelioma. *J Thorac Oncol* 2016; **11**(2):249-255.
- (144) Zaragoza J, Caille A, Beneton N, Bens G, Christiann F, Maillard H et al. High neutrophil to lymphocyte ratio measured before starting ipilimumab treatment is associated with reduced overall survival in patients with melanoma. *Br J Dermatol* 2016; **174**(1):146-151.
- (145) Oki Y, Yamamoto K, Kato H, Kuwatsuka Y, Taji H, Kagami Y et al. Low absolute lymphocyte count is a poor prognostic marker in patients with diffuse large B-cell lymphoma and suggests patients' survival benefit from rituximab. *Eur J Haematol* 2008; **81**(6):448-453.
- (146) Furukawa K, Uwagawa T, Iwase R, Haruki K, Fujiwara Y, Gocho T et al. Prognostic factors of unresectable pancreatic cancer treated with nafamostat mesilate combined with gemcitabine chemotherapy. *Anticancer Res* 2012; **32**(11):5121-5126.
- (147) Lin GN, Peng JW, Xiao JJ, Liu DY, Xia ZJ. Prognostic impact of circulating monocytes and lymphocyte-to-monocyte ratio on previously untreated metastatic non-small cell lung cancer patients receiving platinum-based doublet. *Med Oncol* 2014; **31**(7):70.
- (148) Lin GN, Peng JW, Liu DY, Xiao JJ, Chen YQ, Chen XQ. Increased lymphocyte to monocyte ratio is associated with better prognosis in patients with newly diagnosed metastatic nasopharyngeal carcinoma receiving chemotherapy. *Tumour Biol* 2014; **35**(11):10849-10854.
- (149) Lin GN, Liu PP, Liu DY, Peng JW, Xiao JJ, Xia ZJ. Prognostic significance of the pre-chemotherapy lymphocyte-to-monocyte ratio in patients with previously untreated metastatic colorectal cancer receiving FOLFOX chemotherapy. *Chin J Cancer* 2016; **35**:5.
- (150) Wu ES, Oduyebo T, Cobb LP, Cholakian D, Kong X, Fader AN et al. Lymphopenia and its association with survival in patients with locally advanced cervical cancer. *Gynecol Oncol* 2016; **140**(1):76-82.
- (151) Bari A, Tadmor T, Sacchi S, Marcheselli L, Liardo EV, Pozzi S et al. Monocytosis has adverse prognostic significance and impacts survival in patients with T-cell lymphomas. *Leuk Res* 2013; **37**(6):619-623.
- (152) Cho KS, Choi YD, Kim SJ, Kim CI, Chung BH, Seong dH et al. A comprehensive prognostic stratification for patients with metastatic renal clear cell carcinoma. *Yonsei Med J* 2008; **49**(3):451-458.
- (153) Chen YP, Zhao BC, Chen C, Shen LJ, Gao J, Mai ZY et al. Pretreatment platelet count improves the prognostic performance of the TNM staging system and aids in planning therapeutic regimens for nasopharyngeal carcinoma: a single-institutional study of 2,626 patients. *Chin J Cancer* 2015; **34**(3):137-146.

- (154) Hong X, Cui B, Wang M, Yang Z, Wang L, Xu Q. Systemic Immune-inflammation Index, Based on Platelet Counts and Neutrophil-Lymphocyte Ratio, Is Useful for Predicting Prognosis in Small Cell Lung Cancer. *Tohoku J Exp Med* 2015; **236**(4):297-304.
- (155) Shoultz-Henley S, Garden AS, Mohamed AS, Sheu T, Kroll MH, Rosenthal DI et al. Prognostic value of pretherapy platelet elevation in oropharyngeal cancer patients treated with chemoradiation. *Int J Cancer* 2016; **138**(5):1290-1297.
- (156) Forrest LM, McMillan DC, McArdle CS, Angerson WJ, Dunlop DJ. Evaluation of cumulative prognostic scores based on the systemic inflammatory response in patients with inoperable non-small-cell lung cancer. *Br J Cancer* 2003; **89**(6):1028-1030.
- (157) Elahi MM, McMillan DC, McArdle CS, Angerson WJ, Sattar N. Score based on hypoalbuminemia and elevated C-reactive protein predicts survival in patients with advanced gastrointestinal cancer. *Nutr Cancer* 2004; **48**(2):171-173.
- (158) Crumley AB, McMillan DC, McKernan M, McDonald AC, Stuart RC. Evaluation of an inflammation-based prognostic score in patients with inoperable gastro-oesophageal cancer. *Br J Cancer* 2006; **94**(5):637-641.
- (159) Glen P, Jamieson NB, McMillan DC, Carter R, Imrie CW, McKay CJ. Evaluation of an inflammation-based prognostic score in patients with inoperable pancreatic cancer. *Pancreatol* 2006; **6**(5):450-453.
- (160) Read JA, Choy ST, Beale PJ, Clarke SJ. Evaluation of nutritional and inflammatory status of advanced colorectal cancer patients and its correlation with survival. *Nutr Cancer* 2006; **55**(1):78-85.
- (161) Leitch EF, Chakrabarti M, Crozier JE, McKee RF, Anderson JH, Horgan PG et al. Comparison of the prognostic value of selected markers of the systemic inflammatory response in patients with colorectal cancer. *Br J Cancer* 2007; **97**(9):1266-1270.
- (162) Crumley AB, Stuart RC, McKernan M, McDonald AC, McMillan DC. Comparison of an inflammation-based prognostic score (GPS) with performance status (ECOG-ps) in patients receiving palliative chemotherapy for gastroesophageal cancer. *J Gastroenterol Hepatol* 2008; **23**(8 Pt 2):e325-e329.
- (163) Ramsey S, Aitchison M, Graham J, McMillan DC. The longitudinal relationship between the systemic inflammatory response, circulating T-lymphocytes, interleukin-6 and -10 in patients undergoing immunotherapy for metastatic renal cancer. *BJU Int* 2008; **102**(1):125-129.
- (164) Ishizuka M, Nagata H, Takagi K, Kubota K. Influence of inflammation-based prognostic score on mortality of patients undergoing chemotherapy for far advanced or recurrent unresectable colorectal cancer. *Ann Surg* 2009; **250**(2):268-272.
- (165) Hwang JE, Kim HN, Kim DE, Choi HJ, Jung SH, Shim HJ et al. Prognostic significance of a systemic inflammatory response in patients receiving first-line palliative chemotherapy for recurrent or metastatic gastric cancer. *BMC Cancer* 2011; **11**:489.

- (166) Chua W, Clarke SJ, Charles KA. Systemic inflammation and prediction of chemotherapy outcomes in patients receiving docetaxel for advanced cancer. *Support Care Cancer* 2012; **20**(8):1869-1874.
- (167) Inoue Y, Iwata T, Okugawa Y, Kawamoto A, Hiro J, Toiyama Y et al. Prognostic significance of a systemic inflammatory response in patients undergoing multimodality therapy for advanced colorectal cancer. *Oncology* 2013; **84**(2):100-107.
- (168) Leung EY, Scott HR, McMillan DC. Clinical utility of the pretreatment glasgow prognostic score in patients with advanced inoperable non-small cell lung cancer. *J Thorac Oncol* 2012; **7**(4):655-662.
- (169) Jeong JH, Lim SM, Yun JY, Rhee GW, Lim JY, Cho JY et al. Comparison of two inflammation-based prognostic scores in patients with unresectable advanced gastric cancer. *Oncology* 2012; **83**(5):292-299.
- (170) Gioulbasanis I, Pallis A, Vlachostergios PJ, Xyrafas A, Giannousi Z, Perdikouri IE et al. The Glasgow Prognostic Score (GPS) predicts toxicity and efficacy in platinum-based treated patients with metastatic lung cancer. *Lung Cancer* 2012; **77**(2):383-388.
- (171) Hwang EC, Hwang IS, Yu HS, Kim SO, Jung SI, Hwang JE et al. Utility of inflammation-based prognostic scoring in patients given systemic chemotherapy first-line for advanced inoperable bladder cancer. *Jpn J Clin Oncol* 2012; **42**(10):955-960.
- (172) Laird BJ, Kaasa S, McMillan DC, Fallon MT, Hjerstad MJ, Fayers P et al. Prognostic factors in patients with advanced cancer: a comparison of clinicopathological factors and the development of an inflammation-based prognostic system. *Clin Cancer Res* 2013; **19**(19):5456-5464.
- (173) Linton A, Pond G, Clarke S, Vardy J, Galsky M, Sonpavde G. Glasgow prognostic score as a prognostic factor in metastatic castration-resistant prostate cancer treated with docetaxel-based chemotherapy. *Clin Genitourin Cancer* 2013; **11**(4):423-430.
- (174) Sachlova M, Majek O, Tucek S. Prognostic value of scores based on malnutrition or systemic inflammatory response in patients with metastatic or recurrent gastric cancer. *Nutr Cancer* 2014; **66**(8):1362-1370.
- (175) Zhang P, Xi M, Li QQ, He LR, Liu SL, Zhao L et al. The modified glasgow prognostic score is an independent prognostic factor in patients with inoperable thoracic esophageal squamous cell carcinoma undergoing chemoradiotherapy. *J Cancer* 2014; **5**(8):689-695.
- (176) Anshushaug M, Gynnild MA, Kaasa S, Kvikstad A, Gronberg BH. Characterization of patients receiving palliative chemo- and radiotherapy during end of life at a regional cancer center in Norway. *Acta Oncol* 2015; **54**(3):395-402.
- (177) Moriwaki T, Ishige K, Araki M, Yoshida S, Nishi M, Sato M et al. Glasgow Prognostic Score predicts poor prognosis among advanced biliary tract cancer patients with good performance status. *Med Oncol* 2014; **31**(11):287.
- (178) Miura T, Matsumoto Y, Hama T, Amano K, Tei Y, Kikuchi A et al. Glasgow prognostic score predicts prognosis for cancer patients in palliative settings: a subanalysis of the



Japan-prognostic assessment tools validation (J-ProVal) study. *Support Care Cancer* 2015; **23**(11):3149-3156.

- (179) de Paula PN, Paiva BS, Hui D, Paiva CE. Validation of the Modified Glasgow Prognostic Score in Advanced Cancer Patients Receiving Palliative Care. *J Pain Symptom Manage* 2016; **51**(2):270-277.
- (180) Tan CS, Read JA, Phan VH, Beale PJ, Peat JK, Clarke SJ. The relationship between nutritional status, inflammatory markers and survival in patients with advanced cancer: a prospective cohort study. *Support Care Cancer* 2015; **23**(2):385-391.
- (181) Jung SH, Yang DH, Ahn JS, Kim YK, Kim HJ, Lee JJ. Serum lactate dehydrogenase with a systemic inflammation score is useful for predicting response and survival in patients with newly diagnosed diffuse large B-cell lymphoma. *Acta Haematol* 2015; **133**(1):10-17.
- (182) Xiao Y, Ren YK, Cheng HJ, Wang L, Luo SX. Modified Glasgow prognostic score is an independent prognostic factor in patients with cervical cancer undergoing chemoradiotherapy. *Int J Clin Exp Pathol* 2015; **8**(5):5273-5281.
- (183) Kasuga A, Okano N, Naruge D, Kitamura H, Takasu A, Nagashima F et al. Retrospective analysis of fixed dose rate infusion of gemcitabine and S-1 combination therapy (FGS) as salvage chemotherapy in patients with gemcitabine-refractory advanced pancreatic cancer: inflammation-based prognostic score predicts survival. *Cancer Chemother Pharmacol* 2015; **75**(3):457-464.
- (184) Simmons CP, Koinis F, Fallon MT, Fearon KC, Bowden J, Solheim TS et al. Prognosis in advanced lung cancer--A prospective study examining key clinicopathological factors. *Lung Cancer* 2015; **88**(3):304-309.
- (185) Zhou T, Hong S, Hu Z, Hou X, Huang Y, Zhao H et al. A systemic inflammation-based prognostic scores (mGPS) predicts overall survival of patients with small-cell lung cancer. *Tumour Biol* 2015; **36**(1):337-343.
- (186) Chou WC, Kao CY, Wang PN, Chang H, Wang HM, Chang PH et al. The application of the Palliative Prognostic Index, charlson comorbidity index, and Glasgow Prognostic Score in predicting the life expectancy of patients with hematologic malignancies under palliative care. *BMC Palliat Care* 2015; **14**:18.
- (187) Jiang AG, Chen HL, Lu HY. The relationship between Glasgow Prognostic Score and serum tumor markers in patients with advanced non-small cell lung cancer. *BMC Cancer* 2015; **15**:386.
- (188) Dreanic J, Dhooge M, Barret M, Brezault C, Mir O, Chaussade S et al. Anti-epidermal or anti-vascular endothelial growth factor as first-line metastatic colorectal cancer in modified Glasgow prognostic score 2' patients. *J Cachexia Sarcopenia Muscle* 2015; **6**(3):231-236.
- (189) Song A, Eo W, Lee S. Comparison of selected inflammation-based prognostic markers in relapsed or refractory metastatic colorectal cancer patients. *World J Gastroenterol* 2015; **21**(43):12410-12420.

- (190) Zhou DS, Xu L, Luo YL, He FY, Huang JT, Zhang YJ et al. Inflammation scores predict survival for hepatitis B virus-related hepatocellular carcinoma patients after transarterial chemoembolization. *World J Gastroenterol* 2015; **21**(18):5582-5590.
- (191) Namikawa T, Munekage E, Munekage M, Maeda H, Yatabe T, Kitagawa H et al. Evaluation of Systemic Inflammatory Response Biomarkers in Patients Receiving Chemotherapy for Unresectable and Recurrent Advanced Gastric Cancer. *Oncology* 2016; **90**(6):321-326.
- (192) Arigami T, Uenosono Y, Ishigami S, Okubo K, Kijima T, Yanagita S et al. A Novel Scoring System Based on Fibrinogen and the Neutrophil-Lymphocyte Ratio as a Predictor of Chemotherapy Response and Prognosis in Patients with Advanced Gastric Cancer. *Oncology* 2016; **90**(4):186-192.
- (193) Hsieh MC, Wang SH, Chuah SK, Lin YH, Lan J, Rau KM. A Prognostic Model Using Inflammation- and Nutrition-Based Scores in Patients With Metastatic Gastric Adenocarcinoma Treated With Chemotherapy. *Medicine (Baltimore)* 2016; **95**(17):e3504.
- (194) Yamanaka T, Matsumoto S, Teramukai S, Ishiwata R, Nagai Y, Fukushima M. The baseline ratio of neutrophils to lymphocytes is associated with patient prognosis in advanced gastric cancer. *Oncology* 2007; **73**(3-4):215-220.
- (195) Teramukai S, Kitano T, Kishida Y, Kawahara M, Kubota K, Komuta K et al. Pretreatment neutrophil count as an independent prognostic factor in advanced non-small-cell lung cancer: an analysis of Japan Multinational Trial Organisation LC00-03. *Eur J Cancer* 2009; **45**(11):1950-1958.
- (196) Kao SC, Pavlakis N, Harvie R, Vardy JL, Boyer MJ, van ZN et al. High blood neutrophil-to-lymphocyte ratio is an indicator of poor prognosis in malignant mesothelioma patients undergoing systemic therapy. *Clin Cancer Res* 2010; **16**(23):5805-5813.
- (197) An X, Ding PR, Li YH, Wang FH, Shi YX, Wang ZQ et al. Elevated neutrophil to lymphocyte ratio predicts survival in advanced pancreatic cancer. *Biomarkers* 2010; **15**(6):516-522.
- (198) An X, Ding PR, Wang FH, Jiang WQ, Li YH. Elevated neutrophil to lymphocyte ratio predicts poor prognosis in nasopharyngeal carcinoma. *Tumour Biol* 2011; **32**(2):317-324.
- (199) Chua W, Charles KA, Baracos VE, Clarke SJ. Neutrophil/lymphocyte ratio predicts chemotherapy outcomes in patients with advanced colorectal cancer. *Br J Cancer* 2011; **104**(8):1288-1295.
- (200) Wang S, Zhang Z, Fang F, Gao X, Sun W, Liu H. The neutrophil/lymphocyte ratio is an independent prognostic indicator in patients with bone metastasis. *Oncol Lett* 2011; **2**(4):735-740.
- (201) Lee Y, Kim SH, Han JY, Kim HT, Yun T, Lee JS. Early neutrophil-to-lymphocyte ratio reduction as a surrogate marker of prognosis in never smokers with advanced lung adenocarcinoma receiving gefitinib or standard chemotherapy as first-line therapy. *J Cancer Res Clin Oncol* 2012; **138**(12):2009-2016.

- (202) Kaneko M, Nozawa H, Sasaki K, Hongo K, Hiyoshi M, Tada N et al. Elevated neutrophil to lymphocyte ratio predicts poor prognosis in advanced colorectal cancer patients receiving oxaliplatin-based chemotherapy. *Oncology* 2012; **82**(5):261-268.
- (203) Pinato DJ, Stebbing J, Ishizuka M, Khan SA, Wasan HS, North BV et al. A novel and validated prognostic index in hepatocellular carcinoma: the inflammation based index (IBI). *J Hepatol* 2012; **57**(5):1013-1020.
- (204) He W, Yin C, Guo G, Jiang C, Wang F, Qiu H et al. Initial neutrophil lymphocyte ratio is superior to platelet lymphocyte ratio as an adverse prognostic and predictive factor in metastatic colorectal cancer. *Med Oncol* 2013; **30**(1):439.
- (205) Unal D, Eroglu C, Kurtul N, Oguz A, Tasdemir A. Are neutrophil/lymphocyte and platelet/lymphocyte rates in patients with non-small cell lung cancer associated with treatment response and prognosis? *Asian Pac J Cancer Prev* 2013; **14**(9):5237-5242.
- (206) Yao Y, Yuan D, Liu H, Gu X, Song Y. Pretreatment neutrophil to lymphocyte ratio is associated with response to therapy and prognosis of advanced non-small cell lung cancer patients treated with first-line platinum-based chemotherapy. *Cancer Immunol Immunother* 2013; **62**(3):471-479.
- (207) Fox P, Hudson M, Brown C, Lord S, Gebiski V, De SP et al. Markers of systemic inflammation predict survival in patients with advanced renal cell cancer. *Br J Cancer* 2013; **109**(1):147-153.
- (208) Cetin B, Berk V, Kaplan MA, Afsar B, Tufan G, Ozkan M et al. Is the pretreatment neutrophil to lymphocyte ratio an important prognostic parameter in patients with metastatic renal cell carcinoma? *Clin Genitourin Cancer* 2013; **11**(2):141-148.
- (209) Jafri SH, Shi R, Mills G. Advance lung cancer inflammation index (ALI) at diagnosis is a prognostic marker in patients with metastatic non-small cell lung cancer (NSCLC): a retrospective review. *BMC Cancer* 2013; **13**:158.
- (210) Troppan K, Deutsch A, Gerger A, Stojakovic T, Beham-Schmid C, Wenzl K et al. The derived neutrophil to lymphocyte ratio is an independent prognostic factor in patients with diffuse large B-cell lymphoma. *Br J Cancer* 2014; **110**(2):369-374.
- (211) Kang MH, Go SI, Song HN, Lee A, Kim SH, Kang JH et al. The prognostic impact of the neutrophil-to-lymphocyte ratio in patients with small-cell lung cancer. *Br J Cancer* 2014; **111**(3):452-460.
- (212) Cho IR, Park JC, Park CH, Jo JH, Lee HJ, Kim S et al. Pre-treatment neutrophil to lymphocyte ratio as a prognostic marker to predict chemotherapeutic response and survival outcomes in metastatic advanced gastric cancer. *Gastric Cancer* 2014; **17**(4):703-710.
- (213) Templeton AJ, Pezaro C, Omlin A, McNamara MG, Leibowitz-Amit R, Vera-Badillo FE et al. Simple prognostic score for metastatic castration-resistant prostate cancer with incorporation of neutrophil-to-lymphocyte ratio. *Cancer* 2014; **120**(21):3346-3352.
- (214) Nuhn P, Vaghasia AM, Goyal J, Zhou XC, Carducci MA, Eisenberger MA et al. Association of pretreatment neutrophil-to-lymphocyte ratio (NLR) and overall survival

(OS) in patients with metastatic castration-resistant prostate cancer (mCRPC) treated with first-line docetaxel. *BJU Int* 2014; **114**(6b):E11-E17.

- (215) Sonpavde G, Pond GR, Armstrong AJ, Clarke SJ, Vardy JL, Templeton AJ et al. Prognostic impact of the neutrophil-to-lymphocyte ratio in men with metastatic castration-resistant prostate cancer. *Clin Genitourin Cancer* 2014; **12**(5):317-324.
- (216) Keizman D, Gottfried M, Ish-Shalom M, Maimon N, Peer A, Neumann A et al. Active smoking may negatively affect response rate, progression-free survival, and overall survival of patients with metastatic renal cell carcinoma treated with sunitinib. *Oncologist* 2014; **19**(1):51-60.
- (217) Li X, Chen ZH, Ma XK, Chen J, Wu DH, Lin Q et al. Neutrophil-to-lymphocyte ratio acts as a prognostic factor for patients with advanced hepatocellular carcinoma. *Tumour Biol* 2014; **35**(11):11057-11063.
- (218) Kacan T, Babacan NA, Seker M, Yucel B, Bahceci A, Eren AA et al. Could the neutrophil to lymphocyte ratio be a poor prognostic factor for non small cell lung cancers? *Asian Pac J Cancer Prev* 2014; **15**(5):2089-2094.
- (219) Lin GN, Peng JW, Liu PP, Liu DY, Xiao JJ, Chen XQ. Elevated neutrophil-to-lymphocyte ratio predicts poor outcome in patients with advanced non-small-cell lung cancer receiving first-line gefitinib or erlotinib treatment. *Asia Pac J Clin Oncol* 2014.
- (220) Yoo EJ, Park JC, Kim EH, Park CH, Shim CN, Lee HJ et al. Prognostic value of neutrophil-to-lymphocyte ratio in patients treated with concurrent chemoradiotherapy for locally advanced oesophageal cancer. *Dig Liver Dis* 2014; **46**(9):846-853.
- (221) Langsenlehner T, Thurner EM, Krenn-Pilko S, Langsenlehner U, Stojakovic T, Gerger A et al. Validation of the neutrophil-to-lymphocyte ratio as a prognostic factor in a cohort of European prostate cancer patients. *World J Urol* 2015; **33**(11):1661-1667.
- (222) Jiang L, Jiang S, Situ D, Lin Y, Yang H, Li Y et al. Prognostic value of monocyte and neutrophils to lymphocytes ratio in patients with metastatic soft tissue sarcoma. *Oncotarget* 2015; **6**(11):9542-9550.
- (223) Lorente D, Mateo J, Templeton AJ, Zafeiriou Z, Bianchini D, Ferraldeschi R et al. Baseline neutrophil-lymphocyte ratio (NLR) is associated with survival and response to treatment with second-line chemotherapy for advanced prostate cancer independent of baseline steroid use. *Ann Oncol* 2015; **26**(4):750-755.
- (224) Luo G, Guo M, Liu Z, Xiao Z, Jin K, Long J et al. Blood neutrophil-lymphocyte ratio predicts survival in patients with advanced pancreatic cancer treated with chemotherapy. *Ann Surg Oncol* 2015; **22**(2):670-676.
- (225) Chen ZY, Raghav K, Lieu CH, Jiang ZQ, Eng C, Vauthey JN et al. Cytokine profile and prognostic significance of high neutrophil-lymphocyte ratio in colorectal cancer. *Br J Cancer* 2015; **112**(6):1088-1097.
- (226) Santoni M, Buti S, Conti A, Porta C, Procopio G, Sternberg CN et al. Prognostic significance of host immune status in patients with late relapsing renal cell carcinoma treated with targeted therapy. *Target Oncol* 2015; **10**(4):517-522.

- (227) Ho CL, Lu CS, Chen JH, Chen YG, Huang TC, Wu YY. Neutrophil/Lymphocyte Ratio, Lymphocyte/Monocyte Ratio, and Absolute Lymphocyte Count/Absolute Monocyte Count Prognostic Score in Diffuse Large B-Cell Lymphoma: Useful Prognostic Tools in the Rituximab Era. *Medicine (Baltimore)* 2015; **94**(24):e993.
- (228) Mitchell P, Thatcher N, Socinski MA, Wasilewska-Tesluk E, Horwood K, Szczesna A et al. Tecemotide in unresectable stage III non-small-cell lung cancer in the phase III START study: updated overall survival and biomarker analyses. *Ann Oncol* 2015; **26**(6):1134-1142.
- (229) Wang YY, Bai ZL, He JL, Yang Y, Zhao R, Hai P et al. Prognostic Value of Neutrophil-Related Factors in Locally Advanced Cervical Squamous Cell Carcinoma Patients Treated with Cisplatin-Based Concurrent Chemoradiotherapy. *Dis Markers* 2016; **2016**:3740794.
- (230) Beltran BE, Aguilar C, Quinones P, Morales D, Chavez JC, Sotomayor EM et al. The neutrophil-to-lymphocyte ratio is an independent prognostic factor in patients with peripheral T-cell lymphoma, unspecified. *Leuk Lymphoma* 2016; **57**(1):58-62.
- (231) Zhang GM, Zhu Y, Gu WJ, Zhang HL, Shi GH, Ye DW. Pretreatment neutrophil-to-lymphocyte ratio predicts prognosis in patients with metastatic renal cell carcinoma receiving targeted therapy. *Int J Clin Oncol* 2016; **21**(2):373-378.
- (232) Lee BS, Lee SH, Son JH, Jang DK, Chung KH, Lee YS et al. Neutrophil-lymphocyte ratio predicts survival in patients with advanced cholangiocarcinoma on chemotherapy. *Cancer Immunol Immunother* 2016; **65**(2):141-150.
- (233) Li ZM, Peng YF, Du CZ, Gu J. Colon cancer with unresectable synchronous metastases: the AAAP scoring system for predicting the outcome after primary tumour resection. *Colorectal Dis* 2016; **18**(3):255-263.
- (234) Li J, Jiang R, Liu WS, Liu Q, Xu M, Feng QS et al. A large cohort study reveals the association of elevated peripheral blood lymphocyte-to-monocyte ratio with favorable prognosis in nasopharyngeal carcinoma. *PLoS One* 2013; **8**(12):e83069.
- (235) Rambaldi A, Boschini C, Gritti G, Delaini F, Oldani E, Rossi A et al. The lymphocyte to monocyte ratio improves the IPI-risk definition of diffuse large B-cell lymphoma when rituximab is added to chemotherapy. *Am J Hematol* 2013; **88**(12):1062-1067.
- (236) Go SI, Kim RB, Song HN, Kang MH, Lee US, Choi HJ et al. Prognostic significance of the lymphocyte-to-monocyte ratio in patients with small cell lung cancer. *Med Oncol* 2014; **31**(12):323.
- (237) Koh YW, Jung SJ, Yoon DH, Suh C, Cha HJ, Go H et al. The absolute lymphocyte to monocyte ratio is associated with poor prognosis in classical Hodgkin lymphoma patients younger than 60 years of age. *Hematol Oncol* 2015; **33**(3):133-140.
- (238) Jiang R, Cai XY, Yang ZH, Yan Y, Zou X, Guo L et al. Elevated peripheral blood lymphocyte-to-monocyte ratio predicts a favorable prognosis in the patients with metastatic nasopharyngeal carcinoma. *Chin J Cancer* 2015; **34**(6):237-246.

- (239) Simon Z, Barna S, Miltenyi Z, Husi K, Magyari F, Jona A et al. Combined prognostic value of absolute lymphocyte/monocyte ratio in peripheral blood and interim PET/CT results in Hodgkin lymphoma. *Int J Hematol* 2016; **103**(1):63-69.
- (240) Liu H, Wu Y, Wang Z, Yao Y, Chen F, Zhang H et al. Pretreatment platelet-to-lymphocyte ratio (PLR) as a predictor of response to first-line platinum-based chemotherapy and prognosis for patients with non-small cell lung cancer. *J Thorac Dis* 2013; **5**(6):783-789.
- (241) Li X, Chen ZH, Xing YF, Wang TT, Wu DH, Wen JY et al. Platelet-to-lymphocyte ratio acts as a prognostic factor for patients with advanced hepatocellular carcinoma. *Tumour Biol* 2015; **36**(4):2263-2269.
- (242) Jiang R, Zou X, Hu W, Fan YY, Yan Y, Zhang MX et al. The elevated pretreatment platelet-to-lymphocyte ratio predicts poor outcome in nasopharyngeal carcinoma patients. *Tumour Biol* 2015; **36**(10):7775-7787.
- (243) Nakamura K, Nishida T, Haruma T, Haraga J, Omichi C, Ogawa C et al. Pretreatment platelet-lymphocyte ratio is an independent predictor of cervical cancer recurrence following concurrent chemoradiation therapy. *Mol Clin Oncol* 2015; **3**(5):1001-1006.
- (244) Langsenlehner T, Pichler M, Thurner EM, Krenn-Pilko S, Stojakovic T, Gerger A et al. Evaluation of the platelet-to-lymphocyte ratio as a prognostic indicator in a European cohort of patients with prostate cancer treated with radiotherapy. *Urol Oncol* 2015; **33**(5):201-216.
- (245) Cannon NA, Meyer J, Iyengar P, Ahn C, Westover KD, Choy H et al. Neutrophil-lymphocyte and platelet-lymphocyte ratios as prognostic factors after stereotactic radiation therapy for early-stage non-small-cell lung cancer. *J Thorac Oncol* 2015; **10**(2):280-285.

#### Figure and Table Legends Section:

**Figure 1:** PRISMA flowchart demonstrating study selection

**Figure 2:** Forrest Plot of Studies investigating the prognostic value of CRP in an unselected cohort of patients with advanced cancer

**Figure 3:** Forrest Plot of Studies investigating the prognostic value of Albumin in an unselected cohort of patients with advanced cancer

**Figure 4:** Forrest Plot of Studies investigating the prognostic value of GPS/mGPS in an unselected cohort of patients with advanced cancer

**Figure 5:** Forrest Plot of Studies investigating the prognostic value of NLR in an unselected cohort of patients with advanced cancer

**Figure 6:** Forrest Plot of Studies investigating the prognostic value of LMR in an unselected cohort of patients with advanced cancer

**Figure 7:** Forrest Plot of Studies investigating the prognostic value of PLR in an unselected cohort of patients with advanced cancer

**Supplementary Table:** Studies investigating the prognostic value of all markers of the SIR in an unselected cohort of patients with advanced cancer

**Supplementary Table: Studies investigating the prognostic value of all markers of the SIR in an unselected cohort of patients with advanced cancer.**

No: CRP	Study	Type of Study	Cancer	Country	Patients (n)	Measure of SIR	Systemic Treatment	Cancer deaths (n)	Overall deaths (n)	Cancer Survival (HR, 95% CI)	Overall survival (HR, 95% CI)	Independent Prognostic Factors
1.	Ueno et al <sup>47</sup> 2000	Retrospective	Prostate	Japan	103	CRP: $\geq 50\text{mg/L}$	Active Chemotherapy	N/A	98	N/A	Multivariate: 3.140 (1.51–6.55) $p < 0.010$	Performance Status, CA19-9
2.	McMillan et al <sup>48</sup> 2001	Retrospective	Multiple	UK	772	CRP (Continuous per 10-fold increase in CRP)	Multiple treatments including platinum chemo and radio	596	671	Multivariate: 2.21 (1.92–2.56) $p < 0.0001$	Multivariate: (Non-cancer survival) 5.48 (3.55–8.46) $p < 0.001$	Age, Albumin
3.	Scott et al <sup>49</sup> 2002	Retrospective	Lung	UK	106	CRP $> 10\text{mg/l}$  CRP $> 100\text{mg/l}$	Palliative chemo with supportive treatment but no mention of either specifically	N/A	106	N/A	Multivariate: $> 10\text{mg/l}$ : 1.78 (1.01–3.15) $p = 0.047$ Multivariate: $> 100\text{mg/l}$ : 1.94 (1.41 – 2.65) $p < 0.001$	Age, Tumour Type, Weight Loss, Karnofsky performance status, fatigue
4.	Bromwich et al <sup>50</sup> 2004	Retrospective	Renal	UK	58	CRP $> 10\text{mg/l}$	$\alpha$ -interferon treatment	N/A	55	N/A	Multivariate: 2.03 (1.09–3.80) $p = 0.026$	CRP Only
5.	Elahi et al <sup>51</sup> 2005	Retrospective	Lymphoma	UK	147	CRP ( $\leq 10/11-100/ > 100\text{mg/L}$ )	No mention of treatment but usually treated with chemo	82	147	Multivariate: 8.18 (4.80–13.95) $p < 0.001$	Multivariate: 2.11 (1.22–3.64) $p < 0.001$	CRP Only
6.	Casamassima et al <sup>52</sup> 2005	Retrospective	Renal	Italy	110	CRP: 8mg/L	IL-2 plus gemcitabine and vinorelbine.	N/A	38	N/A	Multivariate: 4.13 (1.68–10.15) $p = 0.002$	DFI less vs. greater than 12
7.	McArdle et al <sup>53</sup> 2006	Prospective	Prostate	UK	62	CRP $> 10\text{mg/L}$	Androgen Deprivation therapy +/- radio	38	41	Multivariate: 1.97 (0.99–3.92) $p = 0.052$	N/A	PSA

8.	Sawaki et al <sup>54</sup> 2006	Retrospective	Pancreatic	Japan	66	CRP: 10-30mg/L  >30mg/L	Gemcitabine 1 <sup>st</sup> line therapy	N/A	56	N/A	Multivariate: 10–30mg/L: 3.08 (1.18– 8.00) p=0.002  >30mg/L 5.99 (2.33–15.45) p=0.002  N/A	Location, diameter of tumour, Liver Mets
9.	Al Murri et al <sup>55</sup> 2006	Retrospective	Breast	UK	96	CRP>10mg/l	Chemotherapy and endocrine therapy	51	N/A	Multivariate: 2.50 (1.40–4.48) p=0.002	GPS	
10.	Nakach et al <sup>56</sup> 2007	Retrospective	Pancreatic	Japan	74	CRP>50mg/L	Second line palliative chemo	71	74	N/A	Performance Status, Peritoneal Dissemination	
11.	Ramsey et al <sup>57</sup> 2007	Retrospective	Renal Cell	UK	119	CRP: >10mg/L	Active Immunotherapy	102	N/A	Multivariate: 2.85 (1.49–5.45) P = 0.002	MSKCC, MRCCPS, GPS, Calcium, Albumin	
12.	Tanaka et al <sup>58</sup> 2008	Retrospective	Pancreatic	Japan	264	CRP>50mg/L	Single-agent gemcitabine therap y	235	264	N/A (PFS given but not CSS)	Karnofsky performance status, TNM stage, Hb, CA19- 9	
13.	Beer et al <sup>59</sup> 2008	Prospective	Prostate	USA	160	CRP: 8mg/L	Docetaxel-based chemotherapy	N/A	63	N/A	CRP Only	
14.	Papadoniou et al <sup>60</sup> 2008	Retrospective	Pancreatic	Greece	215	CRP-5-15mg/L  >15mg/L	Multiple treatments but all palliative	N/A	215	N/A	Univariate: 5-15mg/l: 8.08 (4.26- 15.26) p<0.001  Univariate: >15mg/l: 18.69 (8.23- 42.40) p<0.001	Tumour location in tail, Lymph node spread, Treatment, Performance status, Weight loss, CEA and Jaundice
15.	Yoshida et al <sup>61</sup> 2008	Retrospective	Muscle- invasive bladder cancer	Japan	88	CRP > 5mg/L	ChRT: External beam radio and two cycles of cisplatin	23	N/A	Multivariate: 1.80 (1.01–2.97) p=0.046	N/A	T-stage



16.	Koch et al <sup>62</sup> 2009	Retrospective	NSCLC	Sweden	289	CRP >10mg/L	Palliative supportive care and platinum based chemo	N/A	272	N/A	Multivariate: 1.50 (1.11-2.02) p<0.010	Stage, Performance Status, Smoking, Alb
17.	Hashimoto et al <sup>63</sup> 2009	Retrospective	Pancreatic	Japan	326	CRP>10mg/L	Gemcitabine treatment with palliative intent	N/A	326	N/A	Multivariate: 0.56 (0.42-0.75) p=0.001  Inverse HR: 1.79 (1.33-2.38)	KPS, Liver Mets, Peritoneal Mets, ALP, LDH
18.	Zacharakis et al <sup>64</sup> 2010	Retrospective	Colorectal	Greece	541	CRP: 5-15mg/l >15mg/l	Combination Chemotherapy	N/A	541	N/A	Multivariate: 5-15mg 1.374 (1.051-1.797) p=0.020  >15mg: 1.483 (1.077-2.040) p=0.016	low Hb, Low Alb, Fatigue, Blood transfusions, Combination Chemo, PS change
19.	Iwasa et al <sup>65</sup> 2010	Retrospective	Gastric cancer	Japan	79	CRP>20mg/L	5-FU based chemo	N/A	79	N/A	Multivariate: 2.03 (1.25-3.31) p<0.01	ECOG, Ascites, Alb,
20.	Falkensammer et al <sup>66</sup> 2010	Retrospective	Renal Cell	Austria	86	CRP: >7mg/L	Active Chemotherapy	N/A	N/A (never given in text just HR)	N/A	Univariate: 2.92 (1.58-5.83) p=0.001	Anaemia, Erythropoietin, LDH, Neopeterin
21.	Masago et al <sup>67</sup> 2010	Retrospective	Lung	Japan	79	CRP>10mg/L	Gefitinib chemotherapy	N/A	60	N/A	Multivariate: 1.48 (1.15-1.95) p=0.0073	EGFR
22.	Shimoda et al <sup>68</sup> 2010	Retrospective	Pancreatic	Japan	83	CRP>10mg/L	50 patients received single-agent treatment with gemcitabine (GEM), 9 patients GEM combined with radiotherapy (GEM+R) and 24 patients had best supportive care (BSC).	N/A	83	N/A	Univariate: CRP: 0.92 (0.67-1.27) p=0.6099  Inverse HR: 1.09 (0.79-1.50)	Albumin
23.	Shinohara et al <sup>69</sup> 2011	Retrospective	Renal Cell	Japan	407	CRP>3mg/L	Multiple treatments including Cytokine 362 77 IFN- $\alpha$ , IL-2, Chemo &	307	323	N/A	Multivariate: 2.1 (1.5-3.0) p<0.001	Time from initial diagnosis to metastasis, Hb, corrected Ca, LDH, Liver

									metastasis, Bone Metastasis, Lymph Node Metastasis			
24.	Yi et al <sup>70</sup> 2011	Retrospective	Pancreatic	Korea	298	CRP>12mg/L	Generally poor outcome	N/A	298 (Not specifically mentioned)	N/A	Multivariate: 1.57: (1.07-2.30) p= 0.021	Metastasis to the liver, Ascites or carcinomatosis, Albumin
25.	Kume et al <sup>71</sup> 2011	Retrospective	Renal Cell	Japan	94	CRP>3mg/l	Palliative chemo	N/A	86	N/A	Multivariate: 2.11 (1.13–3.93) p=0.018	Sarcomatid differentiation, Vertebral Bone Involvement, Extraosseous metastasis, ALP
26.	Lee et al <sup>72</sup> 2011	Prospective	Multiple	Korea	126	CRP≥92mg/L	Palliative symptomatic control and chemotherapy	N/A	36	N/A	Multivariate: 2.44 (1.30- 4.60) p=0.006	Chemotherapy
27.	Byström et al <sup>73</sup> 2012	Retrospective	Colorectal	Sweden	106	CRP>5mg/L	Active Chemotherapy	N/A	60	N/A	Univariate: 1.46 (1.176-1.822) p=0.001  Multivariate: 1.11 (0.86-1.44) p=0.435	TPA, TIMP
28.	Ishioka et al <sup>74</sup> 2012	Retrospective	Urothelial	Japan	223	CRP: Continuous	Palliative Chemo and radiotherapy for half with 45% treated with best supportive care	184	184	N/A	Multivariate: 1.60 (1.19–2.15) p=0.001	Age, ECOG PS≥2, Haemoglobin, Log (LDH) Visceral Metastasis, Lymph Node Metastasis
29.	Prins et al <sup>75</sup> 2012	Retrospective	Prostate	USA	119	CRP: continuous, per each doubling of CRP)	End of life symptom care and palliative chemo	N/A	106	N/A	Multivariate: 1.11 (1.02–1.20) p=0.013	Alkaline phosphatase, Haemoglobin
30.	Zeng et al <sup>76</sup> 2012	Retrospective	Laryngeal	China	57	CRP>8mg/L	Palliative chemo-radiotherapy including platinum	29	N/A	Multivariate: 2.66 (1.22–5.82)	N/A	Tumour site (glottic vs. supraglottic vs.

											p=0.014			subglottic)
31.	Pond et al <sup>77</sup> 2012	Retrospective	Prostate	USA and Canada	116	CRP > 8 mg/L				N/A	108		Multivariate: 1.37 (1.13 – 1.66) p=0.002	PCWG-2 Subtype, Risk groups, Halabi nomogram, Smaletz nomogram
32.	Kinoshita et al <sup>78</sup> 2012	Prospective	HCC	Japan	135	CRP > 10mg/L				N/A	123		Multivariate: 3.31 (1.73–6.32) p<0.001	a-Fetoprotein level, Tumour Numbers, Alb, CRP
33.	Morizane et al <sup>79</sup> 2012	Retrospective	Urothelial	Japan	30	CRP > 10mg/L				21	N/A		N/A	CRP Only
34.	Haas et al <sup>80</sup> 2013	Retrospective	Pancreatic	Germany	291	CRP > 10 mg/L but expressed as Log <sup>80</sup>				N/A	237		Multivariate: 1.32 (1.06–1.63) p=0.011	Stage of Disease, Tumour Grading, KPS, Log
35.	Xia et al <sup>81</sup> 2013	Prospective	Nasopharyngeal	China	335	CRP > 2.46mg/L				37	42		Multivariate: 2.114 (1.10- 4.08) p=0.026	Node classification
36.	Yasuda et al <sup>82</sup> 2013	Retrospective	RCC	Japan	52	CRP ≥ 8mg/l				20	22		Multivariate: 1.79 (1.15–2.86) p=0.0099	Neutrophils
37.	Shirakawa et al <sup>83</sup> 2014	Retrospective	Oesophageal	Japan	163	CRP > 10mg/L				N/A	163		Multivariate: 1.631 (1.119–2.376) p=0.011	Performance status, Number of Mets ≥ 3 versus < 3
38.	Teishima et al <sup>84</sup> 2014	Retrospective	Renal Cell	Japan	140	CRP > 3mg/L				70	73		Multivariate: 3.90 (2.06-7.37) P<0.001	Number of Mets, Prior nephrectomy
39.	Deberne et al <sup>85</sup> 2014	Retrospective	Lung	France	55	CRP > 7mg/L				N/A	50		Univariate: 4.3 (2.38-7.8) p<0.001	Leucocytes, Neutrophils, Hb, Alb, ALP, Corrected Ca

40.	Beuselinck et al <sup>186</sup> 2014	Retrospective	Renal Cell	Belgium	200	CRP>5mg/L	best supportive care some palliative surgery as well						Univariate: 3.17 (2.20-4.68) p<0.001	CRP Only
41.	Xue et al <sup>187</sup> 2014	Retrospective	Pancreatic	Japan	269	CRP<5mg/l	Palliative Chemotherapy	231	N/A	N/A			Multivariate: 0.63 (0.41-0.89) p=0.01 Inverse HR: 1.58 (1.12-2.44)	The status of initially unresectable/rec urrent, Distant Mets, ECOG PS, CA19-9, CEA, LDH
42.	Formica et al <sup>188</sup> 2014	Retrospective	Colorectal	USA	106	CRP (Continuous)	Fluorouracil, irinotecan and bevacizumab	N/A	60	N/A			Multivariate: 1.01 (1.00-1.02) p=0.0138	NLR
43.	Kim et al <sup>189</sup> 2014	Prospective	Multiple	Korea	141	CRP>10mg/L	End of life best supportive care	N/A	141	N/A			Multivariate: 1.64 (1.07-2.52) p=0.023	KPS, Time to terminal cancer<12 months, NLR>5
44.	Xue-Feng et al <sup>190</sup> 2015	Retrospective	Lung	China	127	CRP>10mg/L	Palliative Chemotherapy	N/A	127	N/A			Multivariate: 1.80 (1.19-2.71) p=0.005	CEA, Lymph Node N2
45.	Fiala et al <sup>191</sup> 2015	Retrospective	NSCLC	Czech Rep	595	CRP≥10mg/L	Erlotinib	N/A	395	N/A			Multivariate: 1.63 (1.30-2.03) p<0.001	EGFR Status, Stage, ECOG
46.	Adams et al <sup>192</sup> 2015	Retrospective	Diffuse large B cell lymphoma	Netherla nds	104	CRP>10mg/L	Rituximab, Hydroxydaunorubi cin, Oncovin, and prednisolone (R- CHOP).	N/A	34	N/A			Univariate: 2.60 (1.07-6.30) p=0.036	NCCN-IPI
47.	Ito et al <sup>193</sup> 2015	Retrospective	Prostate	Japan	80	CRP>5mg/L	Docetaxel and active chemotherapy	37	38	N/A			Multivariate: 1.95 (1.33-2.96) p<0.001	Hb

48.	Li et al <sup>84</sup> 2015	Retrospective	Osteosarcoma	China	85	CRP>10mg/L	Active Chemotherapy multiple types	N/A	N/A	N/A	Multivariate: 2.39 (1.22-4.67) p=0.01	Tumour size, poor response to chemo, Metastatic disease
49.	Tang et al <sup>85</sup> 2015	Retrospective	Nasopharyngeal	China	1589	hs-CRP>1.96 mg/L	Chemoradiotherapy with chemo being platinum based	N/A	153	N/A	Multivariate: 1.72 (1.24-2.40) p=0.001	Age, Tumour Stage, BMI, EBV DNA
50.	Thurner et al <sup>86</sup> 2015	Retrospective	Prostate	Austria	261	CRP≥8.6mg/L	Conformal Radiotherapy with ADT therapy	24	59	Multivariate: 4.31 (1.22-15.1) p=0.023	Multivariate: 3.24 (1.84-5.71) p<0.001	PSA (10-20)
51.	Zeng et al <sup>87</sup> 2015	Retrospective	Nasopharyngeal	China	79	CRP>8mg/L	Chemoradiotherapy with platinum based chemo	23	N/A	Multivariate: 3.04 (1.22-7.55) p=0.017	N/A	CRP Only
52.	Xu et al <sup>88</sup> 2015	Retrospective	Prostate	China	135	CRP>10mg/L	Palliative care treatment with no mention of type	N/A	124	N/A	Multivariate: 2.39 (1.56-3.69) p<0.001	Gleason Score
53.	Go et al <sup>89</sup> 2015	Retrospective	Lung	Korea	134	CRP≥19mg/L	Palliative chemo in patients with advanced Lung Ca developing VTE	N/A	N/A (Probability of survival given in months)	N/A	Multivariate: 1.596 (0.888-2.865) p=0.118	Stage, Alb, AMC
54.	Martin et al <sup>90</sup> 2015	Retrospective	Pancreatic	Australia	124	CRP>10mg/L	Chemo for metastatic disease and radio for locally advanced	N/A	114	N/A	Multivariate: 1.42 (0.89-2.01) p=0.15	CA19-9, ALC, ANC, Platelet, NLR, PLR, mGPS, Alb, ECOG
55.	Mitsunaga et al <sup>91</sup> 2015	Retrospective and Prospective	Pancreas	Japan	280 (Prospective: 141)	CRP: Inter: >5-20mg/L and High: >20mg/L	GEM chemotherapy	N/A	280 (141 prospective)	N/A	Retrospective Multivariate: Inter: 1.5 (1.1-2.0) p=0.02 High: 2.6 (1.9-3.6) p<0.01 Prospective Multivariate: Inter: 1.5 (0.8-2.8) p=0.19 High: 4.0 (1.6-10.3) p<0.01	Sex, Age, ECOG-PS, UICC stage, CA 19-9, mGPS, NLR

56.	Kim et al <sup>102</sup> 2015	Retrospective	Pancreatic Ductal Ca	Korea	343 (212 underweight palliative chemo)	CRP>10mg/L	FOLFIRINOX and Gemcitabine based chemo	N/A	343	N/A	Multivariate: Whole Group: 2.313 (1.658-3.228) p<0.001 Palliative Chemo: 2.449 (1.635-3.667) p<0.001	ECOG, Alb, NLR Initial site of Mets, No initial chemotherapy
57.	Yao et al <sup>103</sup> 2015	Retrospective	Prostate	Japan	57	CRP>18mg/L	Docetaxel Chemotherapy	N/A	55	N/A	Multivariate: 1.312 (0.428-4.015) p=0.635	Biopsy Gleason Score, PSA values, NLR
58.	Wu et al <sup>104</sup> 2015	Prospective	Lung	China	366	CRP>10.4mg/L	Combination Chemotherapy	N/A	366	N/A	Multivariate: 1.774 (1.270-2.477) p=0.001	Metastasis, NLR
59.	Middleton et al <sup>105</sup> 2016	Retrospective	Pancreatic Ductal Ca	UK	38	CRP (Continuous)	Combination gemcitabine and capecitabine chemo	N/A	38	N/A	Multivariate: 1.55 (1.00-2.39) p=0.049	Log CA19-9
60.	Casadei et al <sup>106</sup> 2016	Prospective	Metastatic Colorectal Ca	Italy	132	hs-CRP (Continuous)	Combination chemotherapy including bevacizumab	N/A	124	N/A	Univariate: 1.006 (1.004-1.009) p<0.0001	N/A
61.	Sheng et al <sup>107</sup> 2016	Retrospective	NSCLC	China	144	CRP (Relatively High vs. Relatively Low)	Combination Chemotherapy	N/A	144	N/A	Univariate 1.43 (0.83-2.47) p=0.204	Current or ex- smoker, stage, ECOG-PS, PNI
62.	Kou et al <sup>108</sup> 2016	Retrospective	Pancreatic	Japan	306	CRP≥5mg/L	Combination chemotherapy with palliative intent	N/A	249	N/A	Multivariate: 1.24 (0.93-1.65) p=0.15	ECOG PS, Distant Metastasis, Initially unresectable, CEA, CA19-9, NLR
63.	Ahn et al <sup>109</sup> 2016	Retrospective	Multiple Cancer Types	Korea	187	CRP≥8.4mg/L	Best supportive care	N/A	187	N/A	Univariate: 1.37 (1.03-1.82) p=0.028	ECOG PS≥3, High PPI score≥6, hyperbilirubinemia
No: Albumin	Study	Type of Study	Cancer	Country	Patients (n)	Measure of SIR	Systemic Treatment	Cancer deaths (n)	Overall deaths (n)	Cancer Survival (HR, 95% CI)	Overall Survival (HR, 95% CI)	Independent Prognostic Factors
1.	Axdorph et al <sup>110</sup> 2000	Retrospective	Hodgkin's disease	UK	145	Alb<40g/L	Multiple treatments including MOPP chemo and radio	48	57	Multivariate: 2.56 (1.05-6.25) p=0.037	N/A	IL-10, Hb<105g/dL

2.	Vigano et al <sup>111</sup> 2000	Retrospective	Multiple palliative cancers	Canada	227	Alb<35g/L	Symptomatic palliative treatment	N/A	208	N/A	Univariate: 1.9 (1.4-2.8) p<0.01	Weight loss, Lymphocyte, Alk- Phos, Karnofsky Performance status, ECOG CA19-9
3.	Maréchal et al <sup>112</sup> 2007	Retrospective	Pancreatic Cancer	Belgium	99	Alb<35g/L	Gemcitabine based chemo as 2 <sup>nd</sup> line	N/A	90	N/A	Multivariate: 4.06 (1.88- 8.77) p<0.001	
4.	Lam et al <sup>113</sup> 2007	Prospective	Multiple	Hong Kong	170	Alb (No threshold)	Palliative supportive treatment	N/A	167	N/A	Multivariate: 0.95 (0.92-0.98) p=0.001  Inverse HR: 1.05 (1.02- 1.09)	Age, Number of Mets, Karnofsky Performance Status, Edmonton Symptom Assessment System
5.	Ramsey et al <sup>157</sup> 2007	Retrospective	Renal Cell cancer  Metastatic	UK	119	Alb<35g/L	Active Immunotherapy	102	N/A	Multivariate: 2.63 (1.38-5.03) P=0.003	N/A	MSKCC, MRCCPS, GPS, Calcium, CRP
6.	Paralkar et al <sup>114</sup> 2008	Retrospective	NSCLC	USA	172	Albs30g/L	Palliative chemotherapy	N/A	159	N/A	Multivariate: 1.7 (1.11-2.76) p=0.02	ECOG PS, Number of Mets
7.	Ngo et al <sup>115</sup> 2008	Retrospective	B-cell lymphoma	Singapor e	183	Alb<37g/L	CHOP (cyclophosphamide, doxorubicin, vincristine, and prednisolone)	N/A	71 (2-year death rates)	N/A	Multivariate: 2.29 (1.28-4.10) p=0.005	Age, LDH, Stage
8.	Iwasa et al <sup>165</sup> 2010	Retrospective	Disseminat ed gastric cancer	Japan	79	Alb<30mg/L	5-FU based chemo	N/A	79	N/A	Multivariate 1.69 (1.05-2.73) p=0.03	ECOG, Ascites, CRP
9.	Shimoda et al <sup>166</sup> 2010	Retrospective	Pancreatic	Japan	83	Alb<35g/L	50 patients received single-agent treatment with gemcitabine (GEM), 9 patients GEM combined with radiotherapy (GEM+R) and 24 patients had best supportive care (BSC).	N/A	83	N/A	Univariate: 7.15 (1.08-47.43) P=0.042	Albumin Only

10.	Shim et al <sup>116</sup> 2011	Retrospective	Gastric Cancer	Korea	502	Alb<40g/L	Taxanes and cisplatin as first line. 2 <sup>nd</sup> line oral fluoropyrimidine monotherapy	N/A	502	N/A	Multivariate: 1.82 (1.32-2.53) P < 0.001	ECOG, Histological grade, PFS< 2.7 months
11.	Yi et al <sup>70</sup> 2011	Retrospective	Pancreatic	Korea	298	Alb<35mg/L	Gemcitabine-based chemotherapy	N/A	298	N/A	Multivariate: 1.701 (1.085-2.667) p=0.021	Metastasis to the liver, Ascites or carcinomatosis, CRP
12.	Trédan et al <sup>117</sup> 2011	Prospective	Multiple	France	299	Alb<38 g/l	Patients treated with palliative chemo but no specific mention of the type	N/A	264	N/A	Multivariate: 1.47 (1.02-2.11) p=0.0374	ECOG, IL-6, LDH, Lymphocyte Count, Platelet Count
13.	Lim et al <sup>118</sup> 2012	Prospective	Biliary Tract Cancer	Korea	50	Alb<35g/L	iFAM chemotherapy in advanced biliary cancer	N/A	49	N/A	Multivariate: 2.11 (1.057-4.22) p=0.034	ECOG, Response to chemotherapy
14.	Prakash et al <sup>119</sup> 2012	Retrospective	B-cell lymphoma	India	486	Alb<40g/L	CHOP Chemo and IFRT chemo in resistant disease	N/A	314	N/A	Univariate: 2.36 (1.32-4.22) p=0.004	Elevated LDH, LR: Not attained, Ages≥60, PS (2,3,4), IPI: Intermediate and high risk, Cycles <6, Hb<10
15.	Kang et al <sup>120</sup> 2014	Retrospective	Biliary Tract	Korea	168	Alb<35g/L	Chemotherapy ultimately palliative. Chemo was platinum based	N/A	168	N/A	Multivariate: 2.0 (1.0-3.8) p=0.036	ECOG, Site of Mets
16.	Ulas et al <sup>121</sup> 2014	Retrospective	Lung Cancer	Turkey	462	Alb<30g/L	Platinum based chemotherapy as both 1 <sup>st</sup> and 2 <sup>nd</sup> line treat	N/A	391	N/A	Multivariate: 1.28 (0.98-1.67) p=0.037	LDH, ECOG, Calcium, Liver Mets, Malignant Pleural effusion, Chemotherapy, No of Mets, LPI
17.	Imedio et al <sup>122</sup> 2014	Retrospective	HCC	Spain	62	ALB<35g/L	TACE chemotherapy sorafenib, followed by second line erlotinib	N/A	44	N/A	Multivariate: 2.99 (1.03-8.66) P=0.044	PS, Alcohol ethology
18.	Malik et al <sup>123</sup> 2014	Retrospective	Renal	USA	70	Alb<34g/L	Bio/chemo or combination therapy	N/A	51	N/A	Multivariate: 2.82 (1.04-7.65) p=0.042	Age, Sex, ECOG, Mets, LDH



19.	Tsai et al <sup>124</sup> 2014	Prospective	Multiple	Taiwan	522	Alb<30g/L	Palliative and supportive care	N/A	479	N/A	Multivariate: 1.98 (1.01-3.88) p<0.05	AST
20.	Stenman et al <sup>125</sup> 2014	Retrospective	Renal Cell Cancer	Sweden	84	Alb<30 g/L	Chemotherapy, Radiotherapy and 20% had Metastectomy	N/A	84	N/A	Multivariate: 2.72 (1.22-6.09) p=0.015	Albumin Only
21.	Koo et al <sup>126</sup> 2015	Retrospective	Gastric Cancer	Korea	3888	Alb<33g/L	Palliative Chemotherapy	N/A	3494	N/A	Multivariate: 1.32 (1.22-1.44) p<0.001	ECOG, No gastronomy, Peritoneal, Bone and Liver Mets, Bilirubin, ALP
22.	Xue-Feng et al <sup>127</sup> 2015	Retrospective	Lung	China	127	Alb: Normal vs. Low	Palliative Chemotherapy	N/A	127	N/A	Multivariate: 0.928 (0.531-1.622) p=0.793 Inverse: 1.078 (0.617-1.883)	CRP, CEA, Lymph Node N2
23.	Kao et al <sup>127</sup> 2015	Retrospective	Multiple	USA	143	Alb<34g/L vs. 24mg/L to 33mg/L vs. <24mg/L	Palliative Radiotherapy	N/A	69	N/A	Multivariate: 2.09 (1.25-3.48) p=0.005	ECOG, Number of Active Tumours, Tumour site
24.	Wild et al <sup>128</sup> 2015	Retrospective	Pancreatic Adenocarcinoma	USA	101	Baseline Alb: continuous	Palliative chemoradiation	86	88	N/A	Multivariate: 3.584 (1.832-6.993) p=0.0002	Lymph Node Count, Baseline Bun and platelets both continuous, PTV: continuous
25.	Helissey et al <sup>129</sup> 2015	Retrospective	Breast Cancer	France	56	Alb<35g/L	CirCe01 phase III trial using platinum chemotherapy	N/A	26	N/A	Multivariate: 11.1 (3.6-34) p<0.001	CTC >5, Receptor Status, Performance Status
26.	Narwani et al <sup>130</sup> 2015	Retrospective	Multiple Myeloma	UK	38	Alb<35g/L	Chemo consists of oral cyclophosphamide 500 mg once weekly; thalidomide 100 mg/d	N/A	22	N/A	Multivariate: 9.34 (2.82-30.92) p<0.001	ALC, Age
27.	Go et al <sup>129</sup> 2015	Retrospective	Lung	Korea	134	Alb<35g/L	Palliative chemo in patients with advanced Lung Ca developing VTE	N/A	N/A (Probability of survival given in months)	N/A	Multivariate: 1.92 (1.07-3.44) p=0.029	Stage, AMC

28.	Martin et al <sup>100</sup> 2015	Retrospective	Pancreatic Cancer	Australia	124	Alb<35g/L vs. >35g/L	Chemo for metastatic disease and radio for locally advanced	N/A	114	N/A	Multivariate: 0.47 (0.31-0.72) p <0.001  Inverse: 2.12 (1.39-3.23)	CA19-9, ALC, ANC, Platelet, NLR, PLR, mGPS, ECOG
29.	Kou et al <sup>108</sup> 2016	Retrospective	Pancreatic Cancer	Japan	306	Alb<35g/L	Combination chemotherapy with palliative intent	N/A	249	N/A	Multivariate: 0.80 (0.59-1.09) p=0.15	ECOG PS, Distant Metastasis, Initially unresectable, CEA, CA19-9, NLR
30.	Moon et al <sup>111</sup> 2016	Prospective	Neck Squamous Cell Ca	Korea	153	Alb<33g/L	Combination chemotherapy and chemoradiotherapy	24	27	Multivariate: 3.80 (1.57-9.19) p=0.003	N/A	ECOG 1/0, BMI <18.5/others, NLR
31.	Uemura et al <sup>112</sup> 2016	Retrospective	Prostate	Japan	41	Alb<39g/L	Combination chemotherapy including docetaxel	22	22 (All patients died of prostate Ca)	Multivariate: 3.776 (1.238- 11.516) p=0.02	Multivariate: 3.776 (1.238-11.516) p=0.02	BSI (>1% vs. ≤1%)
32.	Dorajoo et al <sup>113</sup> 2016	Retrospective	Colorectal	Singapor e	482	Alb<35g/L	Combination chemo for Mets after previous resection of primary tumour	N/A	480	N/A	Multivariate: 1.295 (1.039-1.614) p=0.022	Age≥65, Poorly differentiated Ca, Met site: Live r, Lung, Carcinomatosis, Bone, Carcinoembryoni c antigen
33.	Choi et al <sup>114</sup> 2016	Retrospective	Pancreatic	Korea	396	Alb: Decreased	Palliative Chemotherapy	N/A	396	N/A	Univariate: 1.380 (1.098-1.735) p=0.006	ECOG PS, CA19-9
No: White Blood Cells	Study	Type of Study	Cancer	Country	Patients (n)	Measure of SIR	Systemic Treatment	Cancer deaths (n)	Overall deaths (n)	Cancer Survival (HR, 95%CI)	Overall survival (HR, 95%CI)	Independent Prognostic Factors
1.	Mandreka et al <sup>135</sup> 2006	Retrospective	Lung	USA	1053	WCC> (>10.2x10 <sup>9</sup> /L for males and >10.6x10 <sup>9</sup> /L for females Low)	Chemotherapy majority platinum based	N/A	1011	N/A	Multivariate: 1.44 (1.23-1.69) p=0.001	ECOG, Stage, BMI Underweight, High Hb

2.	Ramsey et al <sup>137</sup> 2007	Retrospective	Renal Cell	UK	119	WCC>11x10 <sup>9</sup> /L	Active Immunotherapy	102	N/A	Multivariate: 1.66 (1.17-2.35) P = 0.004	N/A	MSKCC, MRCCPS, GPS, Calcium, CRP, Albumin
3.	Tibaldi et al <sup>138</sup> 2008	Retrospective	Lung	Italy	320	WCC>10 (>10 x 10 <sup>9</sup> /L)	Chemo Active with cisplatin + gemcitabine or gemcitabine alone	N/A	280	N/A	Multivariate: 1.79 (1.37-2.33) p=0.0001	Performance status, Histology, Brain metastasis
4.	Partridge et al <sup>139</sup> 2012	Retrospective	Multiple	UK	101 (GPS 2)	WCC>10x10 <sup>9</sup> /L	Palliative end of life supportive care	N/A	47 (4-week mortality)	N/A	Multivariate: 1.015 (1.004-1.026) p=0.005	mGPS 2, Age, Primary cancer site: Breast
No: Neutrophils	Study	Type of Study	Cancer	Country	Patients (n)	Measure of SIR	Systemic Treatment	Cancer deaths (n)	Overall deaths (n)	Cancer survival (HR, 95%CI)	Overall survival (HR, 95%CI)	Independent Prognostic Factors
1.	Lee et al <sup>138</sup> 2011	Retrospective	Breast	Australia	693	Neut (log scale) above baseline of ≥7x10 <sup>9</sup> /L	Active chemotherapy as part of two trials	N/A	577	N/A	Multivariate: 1.34 (1.11-1.62) p=0.003	ECOG, ER Status, Number of visceral Mets, Age, Alk Phos, Hb
2.	Kawashima et al <sup>139</sup> 2012	Retrospective	Renal Cell	Japan	87	Neut>ULN	Active Chemotherapy	87	N/A	Multivariate: 3.597 (1.046- 12.364) P=0.042	N/A	Serum Sodium, CRP
3.	Deberne et al <sup>140</sup> 2014	Retrospective	Lung	France	55	Neut: >8000 /mm <sup>3</sup>	Multiple treatments including chemotherapy, radiotherapy, and best supportive care	N/A	50	N/A	Univariate: 3.08 (1.36-7) p=0.0001	Leucocytes, Hb, Alb, Alk P, Corrected Ca, CRP
4.	Luo et al <sup>140</sup> 2015	Retrospective	Nasophary ngeal	China	419	Absolute Neutrophil Count (ANC) >4.7x10 <sup>9</sup> /L	Chemotherapy which was active and cisplatin based	180	N/A	Multivariate: 2.780 (1.819- 4.247) p<0.001	N/A	Age, Stage III/IV, ANC, AER
5.	Lacovelli et al <sup>141</sup> 2015	Retrospective	Renal Cell	Italy	281	Neutrophils >ULN Hb<LLN	Does not seem to mention specifics about chemo	N/A	131	N/A	Multivariate: 1.99 (1.21-3.27) p=0.006	Mets at Diagnosis, ECOG, Hb, Liver Mets
6.	Wu et al <sup>142</sup> 2015	Prospective	Lung	China	366	Neutrophil >3.41x10 <sup>9</sup> cells/ ml	Combination Chemotherapy	N/A	366	N/A	Multivariate: 1.020 (0.655-1.586) p=0.931	Metastasis, NLR, CRP

7.	Ferrucci et al <sup>142</sup> 2016	Prospective	Metastatic Melanoma	Italy	720	ANC≥7500	Ipilimumab	N/A	662	N/A	Multivariate: 3.38 (2.62-4.36) p<0.0001	ECOG, Brain Mets, Liver Mets
8.	Bille et al <sup>143</sup> 2016	Retrospective	Pleural Mesothelioma	USA	191	Neutrophils >ULN	First line combination chemotherapy	N/A	191	N/A	Multivariate: 1.27 (0.82-1.99) p=0.29	Platelet count, Performance status, Histological diagnosis
9.	Zaragoza et al <sup>144</sup> 2016	Retrospective	Melanoma	France	58	Neutrophils: continuous  Neutrophils: ≥7.5x10 <sup>9</sup> /L	Chemotherapy including Ipilimumab	N/A	22	N/A	Univariate: Continuous: 1.34 (1.17- 1.53) p<0.0001  ≥7.5x10 <sup>9</sup> /L : 3.28 (1.38- 7.78) p=0.007	LDH IU, Performance Status
No: Lymphocytes	Study	Type of Study	Cancer	Country	Patients (n)	Measure of SIR	Systemic Treatment	Cancer deaths (n)	Overall deaths (n)	Cancer survival (HR,95%CI)	Overall survival (HR,95%CI)	Independent Prognostic Factors
1.	Oki et al <sup>145</sup> 2008	Retrospective	B-cell Lymphoma	Japan	221	ALC<1x10 <sup>9</sup> /L	Chemotherapy including Rituximab	N/A	N/A (percentage range given)	N/A	Multivariate: 2.51 (1.38-4.58) p=0.003	IPI as a linear parameter
2.	Trédan et al <sup>147</sup> 2011	Prospective	Multiple	France	299	lymphocyte count ≤700/μL	Patients treated with palliative chemo but no specific mention of the type	N/A	264	N/A	Multivariate: 1.43 (1.04-1.95) p=0.0268	ECOG, IL-6, LDH, Alb, Platelet Count
3.	Furukawa et al <sup>146</sup> 2012	Retrospective	Pancreatic	Japan	41	Lymph Count>2000/μl	Nafamostat Mesilate Combined with Gemcitabine Chemotherapy	N/A	41	N/A	Multivariate: 24.016 (5.003-115.278) p<0.0001	Jaundice, Ascites, CA19-9
4.	Lin et al <sup>147</sup> 2014	Retrospective	SCLC	China	370	ALC>0.45x10 <sup>9</sup> / L	Platinum based doublet chemotherapy	N/A	370	N/A	Multivariate: 2.039 (1.488-2.795) p<0.001	LMR, Histology, ECOG

5.	Lin et al <sup>148</sup> 2014	Retrospective	Nasopharyngeal	China	281	ALC<2.25x10 <sup>9</sup> /L	Cisplatin based chemotherapy	N/A	255	N/A	Multivariate: 0.59 (0.43-0.81) p=0.001	Age, LMR
6.	Wild et al <sup>12a</sup> 2015	Retrospective	Pancreatic	USA	101	Lymph (<500 vs. ≥500)	Palliative chemoradiation	86	88	N/A	Multivariate: 2.879 (1.531-5.415) p=0.001	Baseline Alb, Baseline Bun and platelets both continuous, PTV: continuous
7.	Bille et al <sup>143</sup> 2016	Retrospective	Pleural Mesothelioma	USA	191	lymphocyte (>1.4 vs. ≤1.4)	First line combination chemotherapy	N/A	191	N/A	Multivariate: 0.78 (0.54-1.12) P=0.17 Inverse HR: 1.282 (0.893-1.852)	Platelet count, Performance status, Histological diagnosis
8.	Lin et al <sup>149</sup> 2016	Retrospective	Metastatic Colorectal	China	488	ALC ≥2.70x10 <sup>9</sup> /L	FOLFOX chemotherapy	N/A	479	N/A	Multivariate: 0.841 (0.676-1.047) p=0.391 Inverse HR: 1.189 (0.955-1.479)	Gender, ECOG Performance, Tumour differentiation, Pre-chemo AMC and LMR
9.	Wu et al <sup>150</sup> 2016	Retrospective	Cervical Cancer	US	71	TLC>1000 cells/mm <sup>3</sup>	Platinum based chemoradiation	N/A	42	N/A	Multivariate: 0.23 (0.05-1.03) p=0.053 Inverse HR: 4.348 (0.971-20)	Stage III disease
10.	Choi et al <sup>154</sup> 2016	Retrospective	Pancreatic	Korea	396	Lymphocytes <2000 cells/mm <sup>3</sup>	Palliative Chemotherapy	N/A	396	N/A	Univariate: 1.410 (1.119-1.777) p=0.004	ECOG PS, CA19-9
11.	Zaragoza et al <sup>154</sup> 2016	Retrospective	Melanoma	France	58	Lymphocytes: continuous	Chemotherapy including ipilimumab	N/A	22	N/A	Univariate: 0.88 (0.50-1.54) p<0.20 Inverse HR: 1.136 (0.649-2)	LDH IU, Performance Status
No: Monocytes	Study	Type of Study	Cancer	Country	Patients (n)	Measure of SIR	Systemic Treatment	Cancer deaths (n)	Overall deaths (n)	Cancer survival (HR, 95%CI)	Overall survival (HR, 95%CI)	Independent Prognostic Factors

1.	Bari et al <sup>151</sup> 2013	Retrospective	T-cell lymphoma	Italy	94	Mono>0.8x10 <sup>9</sup> /L	Active chemo including vincristine	N/A	48	N/A	Multivariate: 2.41, (1.19–4.89) p=0.015	PIT Score, Histopathology
2.	Lin et al <sup>147</sup> 2014	Retrospective	SCLC	China	370	AMC≥0.45x10 <sup>9</sup> /L	Platinum based doublet chemotherapy	N/A	370	N/A	Multivariate: 0.928 (0.686-1.257) P=0.631	LMR, Histology, ECOG
3.	Lin et al <sup>148</sup> 2014	Retrospective	Nasopharyngeal	China	281	AMC≥0.35x10 <sup>9</sup> /L	Cisplatin based chemotherapy	N/A	255	N/A	Multivariate: 1.20 (0.85-1.70) p=0.309	Age, ALC, LMR
4.	Go et al <sup>149</sup> 2015	Retrospective	Lung	Korea	134	AMC≥640 cells/ $\mu$ L AMC= Absolute Mono Count	Palliative chemo in patients with advanced Lung Ca developing VTE	N/A	N/A (Probability of survival given in months)	N/A	Multivariate: 1.994 (1.137-3.498) p=0.016	Stage, Alb,
5.	Lin et al <sup>149</sup> 2016	Retrospective	Metastatic Colorectal	China	488	AMC ≥0.55x10 <sup>9</sup> /L	FOLFOX chemotherapy	N/A	479	N/A	Multivariate: 1.514 (1.204-1.903) p<0.001	Gender, ECOG Performance, Tumour differentiation, Pre-chemo LMR
No: Platelets	Study	Type of Study	Cancer	Country	Patients (n)	Measure of SIR	Systemic Treatment	Cancer deaths (n)	Overall deaths (n)	Cancer survival (HR, 95%CI)	Overall survival (HR, 95%CI)	Independent Prognostic Factors
1.	Cho et al <sup>152</sup> 2008	Retrospective	Renal Cell	Korea	197	Plate>450,000/ $\text{mm}^3$	Immunotherapy (interferon- $\alpha$ , interleukin-2 (IL-2), or a combination thereof with or without 5-fluorouracil	127	129	Multivariate: 1.34 (0.74 – 2.41) p=0.333	N/A	ECOG-PS, N-stage, Sarcomatoid differentiation, Number of Mets
2.	Trédan et al <sup>117</sup> 2011	Prospective	Multiple	France	299	Plate <130g/L	Patients treated with palliative chemo but no specific mention of the type	N/A	264	N/A	Multivariate: 1.70 (1.02-2.81) p=0.0402	ECOG, IL-6, LDH, Lymphocyte Count, Alb

3.	Stenman et al <sup>123</sup> 2014	Retrospective	Renal Cell	Sweden	84	Plate: >360X 10 <sup>9</sup> /L	Chemo, Radio and 20% had Metastectomy	N/A	84	N/A	Multivariate: 1.62 (0.79–3.32) p=0.19	Albumin Only
4.	Chen et al <sup>133</sup> 2015	Retrospective	Nasopharyngeal	China	2626	Plate>300x10 <sup>9</sup> /L	Active radio and chemo or combination	N/A	774	N/A	Multivariate: 1.810 (1.531–2.140) p<0.001	Age, Sex, T-stage, N-stage
5.	Wild et al <sup>128</sup> 2015	Retrospective	Pancreatic	USA	101	Baseline Plate: continuous	Palliative chemoradiation	86	88	N/A	Multivariate: 1.004 (1.001–1.007) p=0.005	Baseline Alb, LN Count, Baseline Bil both continuous, PTV: continuous
6.	Hong et al <sup>154</sup> 2015	Retrospective	Lung Cancer	China	919	PlatesULN	Chemotherapy and radiotherapy	N/A	892	N/A	Multivariate: 1.016 (0.855–1.208) p=0.856	Stage, Response to treatment, LDH
7.	Shoultz-Henley et al <sup>155</sup> 2016	Retrospective	Oropharyngeal	USA	433	Plate: 350x10 <sup>9</sup> /L	Combined chemo and radiotherapy	N/A	Not mentioned only % given	N/A	Multivariate: 1.9 (1.2–2.9) p<0.006	Anaemia, Dahstrom-Sturgis category, HPV status
8.	Bille et al <sup>143</sup> 2016	Retrospective	Pleural Mesothelioma	USA	191	Plate>450,000 per mm <sup>3</sup>	First line combination chemotherapy	N/A	191	N/A	Multivariate: 2.09 (1.33–3.35) p=0.002	Performance status, Histological diagnosis
No: GPS/mGPS	Study	Type of Study	Cancer	Country	Patients (n)	Measure of SIR	Systemic Treatment	Cancer deaths (n)	Overall deaths (n)	Cancer survival (HR, 95%CI)	Overall survival (HR, 95%CI)	Independent Prognostic Factors
1.	Forrest et al <sup>156</sup> 2003	Retrospective	NSCLC	UK	161	GPS (0/1/2)	Chemotherapy mainly cisplatin and radical radio	N/A	118	N/A	Multivariate: 1.111 (1.23–2.35) P= 0.001	Stage/ECOG score, CRP/Alb score
2.	Elahi et al <sup>157</sup> 2004	Retrospective	Gastric and colorectal	UK	165	GPS (0/1/2)	Palliative Chemo and Supportive Care	N/A	165	N/A	Univariate: Gastric: 1.71 (1.15–2.25) P = 0.002 Colorectal: 1.77 (1.51–2.57) P < 0.001)	Age, Tumour Type

3.	Crumley et al <sup>158</sup> 2006	Retrospective	Gastro-oesophageal	UK	258	GPS (0/1/2)	Chemotherapy and radiotherapy with palliative intent	202	211	Multivariate: 1.51 (1.22-1.86) p<0.001	N/A	GPS Only
4.	Al Murri et al <sup>155</sup> 2006	Retrospective	Breast	UK	96	GPS (0/1/2)	Chemotherapy and endocrine therapy	51	N/A	Multivariate: 2.26 (1.45-3.52) p<0.001	N/A	CRP, Alb
5.	Glen et al <sup>159</sup> 2006	Retrospective	Pancreatic	UK	187	GPS (0/1/2)	Palliative treatment with platinum based chemo	N/A	181	N/A	Multivariate: 1.72 (1.40-2.11) p<0.001	Age, TNM
6.	Read et al <sup>160</sup> 2006	Prospective	Colorectal	Australia	51	GPS (0/1/2)	Chemo and Radiotherapy as well as supportive care	N/A	32	N/A	Multivariate: 2.27 (1.09-4.73) P = 0.028	Type of treatment, PS, SAP
7.	Ramsey et al <sup>167</sup> 2007	Retrospective	Renal Cell	UK	119	GPS: (0/1/2)	Active Immunotherapy	102	N/A	Multivariate: 2.35(1.51-3.67) P<0.001	N/A	MSKCC, MRCCPS, Calcium, CRP, Albumin
8.	Leitch et al <sup>161</sup> 2007	Retrospective	Colorectal Liver Mets	UK	84	GPS (0,1,2)	Palliative chemotherapy	71	N/A	Multivariate: 1.44 (1.01-2.04) P = 0.043	N/A	Age, Single liver metastasis, Extra-hepatic disease, chemotherapy treatment
9.	Crumley et al <sup>162</sup> 2008	Retrospective	Gastro-oesophageal	UK	65	GPS (0/1/2)	Mostly cisplatin based chemotherapy	58	59	Multivariate: 1.69 (1.00-2.86) P=0.05	N/A	GPS Only
10.	Ramsey et al <sup>163</sup> 2008	Prospective	Renal Cell	UK	23	GPS (0/1/2)	Palliative treatment with immunotherapy	N/A	15	N/A	Multivariate: 2.23 (1.06-4.57) p=0.029	GPS Only
11.	Ishizuka et al <sup>164</sup> 2010	Retrospective	Colorectal	Japan	112	mGPS: 1/2	Active chemo in form of FOLFIRI and FOLFOX regimens	44	N/A	Multivariate: 6.071 (1.625-22.68) p=0.0073	N/A	mGPS only
12.	Shimoda et al <sup>166</sup> 2010	Retrospective	Pancreatic	Japan	83	GPS (0 vs. 1 or 2)	50 patients received single-agent treatment with gemcitabine (GEM), 9 patients GEM combined with	N/A	83	N/A	Univariate: 0.513 (0.047-5.547) P=0.5825 Inverse: 1.949 (0.180-21.277)	Albumin



[illegible]

21.	Laird et al <sup>172</sup> 2013	Prospective	Multiple	UK	1825 (Test)  631 (Validation n)	GPS: 1&2	Chemo, radio and BSC	N/A	1601(Test)  471 (Validation)	N/A	GPS 2: 7.00 (2.53-19.36) p=0.001	Test: Dyspnoea, ECOG  Validation: Quality of life, Physical Function, Pain, BMI, ECOG
22.	Linton et al <sup>173</sup> 2013	Prospective	Prostate	Australia	112	mGPS (2 vs. 0)  (1 vs. 0)	Docetaxel and prednisone treatment	N/A	84	N/A	Univariate: mGPS Categorical (2 vs. 0) 3.44 (1.75-6.76) p <0.001  (1 vs 0) 1.97 (1.01-3.83) p=0.047	mGPS only
23.	Sachlova et al <sup>174</sup> 2014	Retrospective	Gastric	Czech Rep	64 (treated with chemo)	GPS (1&2)	Palliative chemo mostly platinum based	N/A	64	N/A	Multivariate: GPS 1: 1.93 (0.85-4.40) p=0.12  GPS 2: 6.63 (2.42-18.17) p<0.001	OPNI
24.	Zhang et al <sup>175</sup> 2014	Retrospective	Oesophageal	China	212	mGPS (0,1,2)	Radiotherapy and cisplatin based chemo	N/A	160	N/A	Multivariate: 1.694 (1.350-2.126) p<0.001	Location, T&M, stage
25.	Anshushaug et al <sup>176</sup> 2014	Retrospective	Multiple	Norway	723	GPS (1 & 2)	Palliative radio and chemo	N/A	723	N/A	Multivariate: Chemo: GPS: 1: 1.69 (0.72-4.00) p=0.23  2: 3.00 (1.32-6.80) p=0.009	Age, Performance status, Referred to Palliative Care, Mets when diagnosed

26.	Moriwaki et al <sup>177</sup> 2014	Retrospective	Biliary Tract	Japan	218	Continuous: GPS (0 vs. 1/2)	Chemo with GEM and CDDP regimens	N/A	218	N/A	Radio: GPS 1: 2.90 (0.97-8.67) p=0.06 GPS 2: 3.98 (1.52-10.42) p=0.005  Multivariate: 0.60 (0.40-0.90) P=0.012 Inverse: 1.666 (1.111-2.5)	ALP, LDH, No of Mets, Liver, Peritoneal/Other Mets
27.	Miura et al <sup>178</sup> 2015	Prospective	Multiple	Japan	1160	GPS 1&2	Purely palliative care no active treatment	N/A	1160 (All end of life care)	N/A	Multivariate: GPS 1: 1.07 (0.78-1.49) P= 0.673 GPS 2: 1.36 (1.01-1.87) P= 0.046	Performance status, Liver Mets, PP >6l, NLR24, Dyspnea, Oedem0.308a
28.	De Paula Pantano et al <sup>179</sup> 2015	Prospective	Multiple	USA	459	mGPS 1&2	Predominantly supportive treatment but some still receiving chemo	N/A	346	N/A	Multivariate: GPS 1: 2.066 (1.356-3.147) P= 0.001 GPS 2: 2.664 (1.929-2.680) P<0.001	Sex, Hepatic Mets, CNS Mets, Treatment Palliative care only, KPS (0-70%),
29.	Tan et al <sup>180</sup> 2015	Prospective	Multiple	Australia	114	mGPS: 1/2	Chemotherapy but no specific mention of type	N/A	Followed up until the date of death or the date that data as last updated.	N/A	Multivariate: 1.68 (1.03-2.76) p=0.039	PG-SGA C, Required dose reduction +/- transfusion
30.	Jung et al <sup>181</sup> 2015	Retrospective	B-cell Lymphoma	Korea	213	L-GPS: 1&2	R-CHOP chemotherapy.	50	58	N/A	Multivariate GPS 1: 2.135 (0.919-4.533) p=0.078 GPS 2: 5.898 (2.028-14.454) p=0.001	ECOG
31.	Xiao et al <sup>182</sup> 2015	Retrospective	Cervical	China	238	mGPS (0/1/2)	Chemo in the form of Cisplatin plus 5-fluorouracil or cisplatin plus docetaxel. Also, treated with radio	N/A	124	N/A	Multivariate: 1.820 (1.378-2.404) p<0.001	PS, FIGO Stage, LN status

32.	Martin et al <sup>100</sup> 2015	Retrospective	Pancreatic	Australia	124	mGPS: (0,1,2)	Chemo for metastatic disease and radio for locally advanced	N/A	114	N/A	Multivariate: mGPS 1: 41 (1.10-1.80) p=0.01	CA19-9, ALC, ANC, Platelet, NLR, PLR, Alb, ECOG
33.	Kasuga et al <sup>103</sup> 2015	Retrospective	Pancreatic	Japan	61	mGPS: 2	Gemcitabine and S-1 combination therapy (FGS) as salvage chemotherapy	N/A	61	N/A	Multivariate: 6.605 (2.965-14.709) p<0.001	CA19-9 > 2,000 ECOG>0
34.	Simmons et al <sup>104</sup> 2015	Prospective	Lung	Greece	390	mGPS (0/1/2)	Best supportive care	N/A	283	N/A	Multivariate: 1.67 (1.40-2.00) p<0.001	ECOG
35.	Zhou et al <sup>105</sup> 2015	Retrospective	Lung	China	359	mGPS 1&2	Radiotherapy and chemotherapy (Irinotecan, Etoposide)	N/A	180	N/A	Multivariate: mGPS 1: 1.52 (1.08-2.13) p=0.015  mGPS 2: 5.23 (2.36-11.58) p<0.001	Adjusted for age, sex, disease stage, ECOG-PS.
36.	Chou et al <sup>106</sup> 2015	Retrospective	Haematological	China	217	GPS: (1&2)	Palliative care no specific mention of chemo	N/A	204	N/A	Multivariate: GPS 1: 2.12 (1.13-3.97) p=0.020  GPS 2: 1.71 (0.964-3.05) p=0.069	PPI > 4.5.
37.	Jiang et al <sup>107</sup> 2015	Prospective	Lung	China	138	GPS: 1&2	Cisplatin based chemo.	N/A	138	Multivariate: GPS 1: 0.8 (0.5-0.9) p=0.02  GPS 2: 0.6 (0.2-0.8) p=0.02  Inverse: GPS 1: 1.25 (1.111-2)  GPS 2: 1.666 (1.25-5)	CYFRA21-1, CEA, TPS	
38.	Dreanic et al <sup>108</sup> 2015	Retrospective	Colorectal	France	27	mGPS: 2  Inverse mGPS: 2	5-fluorouracil-based systemic chemotherapy and anti-VEGF	N/A	27	N/A	Univariate in anti-VEGF group: 0.48 (0.18-1.29) p=0.15  Inverse: 2.083 (0.775-	GPS Only

												5.555)		
39.	Mitsunaga et al <sup>101</sup> 2015	Pancreas	Japan	Prospective	280 (Prospective: 141)	mGPS: 1 & 2	GEM chemotherapy	N/A	280 (141 prospective)	N/A	Multivariate: mGPS: 1: 0.9(0.4-1.9) p=0.76  mGPS 2: 0.72 (0.3-1.7) p=0.47  Inverse: mGPS 1: 1.111 (0.526-2.5)  mGPS 2: 1.388 (0.588-3.333)	Sex, Age, ECOG-PS, UICC stage, CA 19-9, Prognostic CRP Classification, NLR		
40.	Song et al <sup>189</sup> 2015	Colorectal	Korea	Retrospective	177	mGPS: (0 vs. 1 or 2)	Best supportive care and herbal therapy	N/A	177	N/A	Multivariate: 0 vs 1: 1.135 (0.717-1.797) p=0.588  0 vs 2: 3.212 (1.437-7.716) p=0.004  Multivariate: 1.697 (1.325-2.174) p<0.001	LMR, CA19-9, AST, KM treatment		
41.	Zhou et al <sup>190</sup> 2015	HCC	China	Prospective	244	GPS (0/1/2)	TRACE chemotherapy	N/A	198	N/A		ALT, CLIP score		
42.	Namikawa et al <sup>191</sup> 2016	Gastric	Japan	Retrospective	224	GPS (0/1 or 2)  mGPS (0/1 or 2)	Combination chemotherapy including trastuzumab	N/A	223	N/A	Multivariate: GPS: 1.297 (0.667-2.552) p=0.444  mGPS: 0.68 (0.350-1.322) p=0.255	Histological type, NLR		
43.	Arigami et al <sup>192</sup> 2016	Gastric	Japan	Retrospective	68	GPS: 1&2	Chemotherapy and chemoradiotherapy	N/A	68	N/A	Multivariate: GPS 1: 0.830 (0.418-1.618) p=0.586  GPS 2: 2.608 (0.792-7.965) p=0.111	F-NLR score (combined fibrinogen and NLR)		
44.	Hsieh et al <sup>193</sup> 2016	Gastric	Taiwan	Retrospective	256	mGPS (>1)	Combination Chemotherapy	N/A	248	N/A	Multivariate: 2.78 (1.60-4.83) p<0.001	Peritoneal Mets, NLR, mGPS, PG-SGA		

No: NLR	Study	Type of Study	Cancer	Country	Patients (n)	Measure of SIR	Systemic Treatment	Cancer deaths (n)	Overall deaths (n)	Cancer survival (HR, 95%CI)	Overall survival (HR, 95%CI)	Independent Prognostic Factors
1.	Yamanaka et al <sup>154</sup> 2008	Prospective	Gastric	Japan	1220	NLR>2.5	Patients receiving oral fluoropyrimidine	N/A	984	N/A	Multivariate: 1.52 (1.32–1.75) p=0.077	Age, ECOG, Advanced Disease, Liver Mets, WBC 9000-12000/mm3
2.	Teramukai et al <sup>155</sup> 2009	Prospective	Lung	Japan	388	NLR≥4.744	Vinorelbine, gemcitabine, docetaxel, paclitaxel, carboplatin	N/A	276	N/A	Multivariate: 1.56 (1.09–2.24) p=0.015	Neutrophil count
3.	Kao et al <sup>156</sup> 2010	Retrospective	Malignant mesothelioma	Australia	173	NLR≥5	Platinum based chemotherapy	N/A	131	N/A	Multivariate: 2.7 (1.8-3.9) p<0.001	Histological subtype
4.	An et al <sup>157</sup> 2010	Retrospective	Pancreatic	China	95	NLR>5	Gemcitabine-based chemotherapy	N/A	95	N/A	Multivariate: 4.489 (1.372–14.692) p=0.013	CA19-9
5.	Chua et al <sup>158</sup> 2011	Prospective	Multiple	Australia	68	NLR>5	Single unit docetaxel treatment	N/A	68	N/A	Multivariate: 2.0 (1.2–3.3) p=0.010	GPS: 1&2
6.	An et al <sup>159</sup> 2011	Prospective	Nasopharyngeal	China	363	NLR>3.73	Local radio and cisplatin and/or 5-FU-based neoadjuvant	96	102	Multivariate: 1.74 (1.15–2.62) p=0.008	N/A	Age, sex and T-stage
7.	Chua et al <sup>159</sup> 2011	Retrospective	Colorectal	Australia	349	NLR≥5	Chemotherapy and best supportive care	N/A	315	N/A	Multivariate: 1.7 (1.2-2.5) p=0.002	ECOG>1
8.	Wang et al <sup>160</sup> 2011	Retrospective	Multiple	China	497	NLR>3	Multiple treatment modalities.	N/A	464	N/A	Multivariate: 1.348 (1.062-1.712) p=0.014	Gender, Tumour Type, Surgery, Other Mets, Adjuvant treatment

9.	Jeong et al <sup>169</sup> 2012	Retrospective	Gastric	Korea	104	NLR>3	Treated with palliative chemo	N/A	94	N/A	Multivariate: 1.65 (1.03–2.64) p = 0.037	Histology, LN Mets, mGPS
10.	Lee et al <sup>201</sup> 2012	Prospective	Lung adenocarcinoma	Korea	199	NLR >2.18	Geftinib with gemcitabine plus cisplatin as first-line therapy.	N/A	N/A (Expressed in months)	N/A	Multivariate: 1.13 (1.06-1.21) p<0.001	ECOG
11.	Kaneko et al <sup>202</sup> 2012	Retrospective	Colorectal	Japan	50	NLR ≥4	Palliative Oxaliplatin-based combination chemotherapy	27	27	N/A	Multivariate: 4.39 (1.82-10.7) p = 0.0013	Platelets
12.	Pinato et al <sup>203</sup> 2012	Retrospective	HCC	USA	112	NLR>5	Active platinum based chemo	N/A	81	N/A	Multivariate: 2.06 (1.16-3.66) p=0.013	IBI, CLIP, BSC
13.	He et al <sup>204</sup> 2012	Retrospective	Colorectal	China	243	NLR≤3 Inverted NLR NLR≥3	Combination chemotherapy including Oxaliplatin and Irinotecan	N/A	199	N/A	Multivariate: 0.678 (0.479-.0961) p=0.029 Inverted: 1.475 (1.041-2.088)	CEA
14.	Linton et al <sup>173</sup> 2013	Prospective	Prostate	Australia	112	NLR: Continuous Categorical: (≥5 vs. <5)	Docetaxel and prednisone treatment	N/A	84	N/A	Univariate: NLR: Cont 1.08 (0.83-1.41) p=0.55 NLR (≥5 vs. <5): 0.98 (0.64-1.49) p=0.91	mGPS 2 vs. 0 and mGPS 1 vs. 0
15.	Unal et al <sup>205</sup> 2013	Prospective	NSCLC	Turkey	94	NLR (low or high)	Chemoradiotherapy including platinum based treat	N/A	81	N/A	Univariate: 1.81 (1.16-2.81) p=0.0008	PLR, Response to chemoradiotherapy
16.	Yao et al <sup>206</sup> 2013	Prospective	Lung	China	182	NLR>2.68	First-line platinum-based chemotherapy.	N/A	91	N/A	Multivariate: 1.761 (1.095–2.832) p=0.020	Nodal spread N2, Metastasis M2.
17.	Fox et al <sup>207</sup> 2013	Retrospective	Renal Cell	Australia	362	NLR>3	Patients treated with Lapatinib or hormonal therapy after prior failure of immunotherapy in a	N/A	357	N/A	Multivariate: 1.42 (1.10- 1.84) p=0.008	Neutrophils, Platelets, KPS, Corrected Calcium, Low Hb

18.	Cetin et al <sup>128</sup> 2013	Retrospective	Renal Cell	Turkey	100				NLR>3.04	Tyrosine Kinase Inhibitors	N/A	54	N/A	Multivariate: 2.406 (1.327-4.361) p=0.004	Male, PFS
19.	Jafri et al <sup>129</sup> 2013	Retrospective	Lung	USA	173				NLR<5	Treated with active chemotherapy multiple types		173		Univariate: 0.57 (0.41-0.79) 0.0008 Inverted HR: 1.754 (1.266-2.439)	PS (0-1/ 2-4), Mets (1-2/ >2), No chemotherapy, ALC <1, ALI < 18
20.	Troppan et al <sup>130</sup> 2014	Retrospective	B-cell Lymphoma	Austria	290				NLR≥4	Standard rituximab, cyclophosphamide, doxorubicin, vincristine, and prednisone (R-CHOP) regimen every 3 weeks for six to eight cycles	N/A	92	N/A	Multivariate: 2.03 (1.17-3.50) p=0.011	Age ≥60, Clinical Stage III & IV, non-GCB, dNLR≥4
21.	Kim et al <sup>131</sup> 2014	Prospective	Multiple	Korea	141				NLR>5	Best supportive care	N/A	141	N/A	Multivariate: 1.96 (1.17-3.31) p=0.011	KPS, Time to terminal cancer<12 months, CRP≥10mg/dl
22.	Kang et al <sup>131</sup> 2014	Retrospective	Lung	United Kingdom	187				NLR≥4	platinum-based chemotherapy	N/A	187	N/A	Multivariate: 1.465 (1.012-2.119) p=0.043	Extensive Disease, LDH
23.	Cho et al <sup>132</sup> 2014	Prospective	Gastric	Korea	268				NLR>3	Initial treatment with chest wall radiotherapy and FOLFOX and platinum based chemo	N/A	268	N/A	Multivariate: 1.569 (1.227-2.006) P<0.001	Undifferentiated, Progressive disease
24.	Templeton et al <sup>133</sup> 2014	Retrospective	Prostate	Canada	357				NLR>3	Docetaxel and platinum based chemo	N/A	345	N/A	Multivariate: 1.89 (1.27-2.82) p=0.002	Liver Mets, Hb, Alb, Log (PSA, LDH, ALP)



25.	Nuhn et al <sup>214</sup> 2014	Retrospective	Prostate	USA	238	NLR>3	First line docetaxel	N/A	237	N/A	Multivariate: 1.883 (1.248, 2.842) p=0.002	Number of chemo cycles, Hb, Alb, AST, Baseline PSA
26.	Sonpavde et al <sup>215</sup> 2014	Retrospective	Prostate	Multinational (US and Canada)	784	NLR (Log transformed)	Patients treated with Sunitinib and prednisolone and docetaxel-based chemotherapy	N/A	516	N/A	Multivariate: 1.55 (1.32-1.83) p<0.001	Log (LDH), Hb, Organ Involvement
27.	Keizmann et al <sup>216</sup> 2014	Prospective	Renal cell	Multinational (USA and Israel)	244	NLR>3	Sunitinib treatment	N/A	203	N/A	Multivariate: 2.95 (2-4.34) p<0.001	Sunitinib induced HTN, Pre-treatment, never having smoked
28.	Li et al <sup>217</sup> 2014	Retrospective	HCC	China	205	NLR>2.43	Sorafenib based chemoembolization	N/A	132	N/A	Multivariate: 1.104 (1.044-1.167) p<0.001	AFP, Tumour Morphology, Child-Pugh Score, Platelets
29.	Formica et al <sup>218</sup> 2014	Retrospective	Colorectal	USA	106	NLR (Continuous)	Fluorouracil, irinotecan and bevacizumab	N/A	60	N/A	Multivariate: 1.8012 (0.2833-1.6048) p=0.0019	CRP
30.	Kacan et al <sup>219</sup> 2014	Retrospective	Lung	Turkey	299	NLR≥5	Chemo and Radiotherapy no mention of surgery	N/A	204 (2 Year survival)	N/A	Multivariate: 1.7 (1.0-2.7) p=0.017	Age, Anaemia at diagnosis, Stage, ECOG PS
31.	Lin et al <sup>219</sup> 2014	Retrospective	Lung	China	81	NLR>3.5	EGFR-TKI treatment	56	56 (All deaths Ca related)	N/A	Multivariate: 3.29 (1.62-6.71) p<0.001	ECOG
32.	Yoo et al <sup>220</sup> 2014	Retrospective	Lung	Korea	138	NLR≥2	Concurrent chemoradiotherapy	N/A	112	N/A	Multivariate: 2.115 (1.193-3.749) p=0.010	ECOG performance status
33.	Langsenlehner et al <sup>221</sup> 2015	Retrospective	Prostate	Austria	415	NLR≥5	Androgen deprivation therapy, Chemotherapy	N/A	60	N/A	Multivariate: 2.16 (1.17-3.99) p=0.013	Intermediate risk group classification

34.	Jiang et al <sup>22</sup> 2015	Retrospective	Soft Tissue Sarcoma	China	154	NLR>1	Treated with active chemotherapy and Ipilimumab	65	80	N/A	Multivariate: 2.477 (1.423-4.311) P= 0.033	Monocyte Ratio>1
35.	Lorente et al <sup>23</sup> 2015	Retrospective	Prostate	United Kingdom	755	NLR: Continuous NLR>3	Patients treated with cabazitaxel (25 mg/m <sup>2</sup> ) versus 3- weekly mitoxantrone (12 mg/m <sup>2</sup> ), both in combination with prednisone 10 mg daily	N/A	N/A (Does not give a figure)	N/A	Multivariate: Conti: 1.91 (1.31-2.79) p=0.001 NLR>3: 1.55 (1.3-1.84), P < 0.001	Measurable disease, Pain at baseline, Treatment arm
36.	Luo et al <sup>24</sup> 2015	Retrospective	Pancreatic	China	403	NLR: ≥3.1	74.9% underwent gemcitabine-based chemotherapy	N/A	394	N/A	Multivariate: 1.42 (1.15-1.74) p=0.001	Age, CA19-9, Albumin, Tumour spread
37.	Kim et al <sup>25</sup> 2015	Retrospective	Pancreatic Ductal	Korea	343 (212 palliative chemo)	NLR>5	FOLFIRINOX and Gemcitabine based chemo	N/A	343	N/A	Multivariate: Whole Group: 1.428 (1.014-2.012) p=0.042 Palliative Chemo: 1.038 (0.654-1.650) p=0.175	ECOG, Alb, CRP, Initial site of Mets, No initial chemotherapy
38.	Chen et al <sup>25</sup> 2015	Retrospective	Colorectal	United States	166	NLR>5	Best supportive care after failure of other treatment in palliative group and Panitumumab in active treatment group	N/A	N/A (No specific numbers of deaths)	N/A	Multivariate: 1.73 (1.03-2.89) p=0.039	Metastatic Site numbers ≥1, LDH>ULN
39.	Santoni et al <sup>26</sup> 2015	Retrospective	Renal Cell	Italy	151	NLR>3	Active treatment with VEGFR-TKI also treated with sunitinib, sorafenib, and pazopanib	N/A	53	N/A	Multivariate: 2.21 (1.21-4.04) p=0.010	MSKCC Prognostic Group
40.	Ho et al <sup>27</sup> 2015	Retrospective	Large B Cell Lymphoma	Taiwan	148	NLR>4.35	Standard R- chemotherapy.	N/A	41	N/A	Multivariate: 1.624 (0.827-3.189) p=0.159	Age, B- symptoms, ECOG, ALC, AMC, ALC/AMC PS

41.	Mitchell et al <sup>128</sup> 2015	Prospective	Lung		Canada	1239	NLR>5		Tecemotide in unresectable stage III non-small-cell lung cancer	N/A	1239	N/A	Univariate: 0.81 (0.66-0.99), P = 0.0383 Inverse HR: 1.235 (1.01-1.515)	High sMUC1, High ANA
42.	Martin et al <sup>100</sup> 2015	Retrospective	Pancreatic		Australia	124	NLR≥5		Chemo for metastatic disease and radio for locally advanced	N/A	114	N/A	Multivariate: 1.60 (1.07-2.40) p=0.02	CA19-9, ALC, ANC, Platelet, PLR, mGPS, Alb, ECOG
43.	Mitsunaga et al <sup>101</sup> 2015	Prospective	Pancreas		Japan	280 (Prospective: 141)	NLR≥5		GEM chemotherapy	N/A	280 (141 prospective)	N/A	Multivariate: 1.3 (0.8-2.2) p=0.32	Sex, Age, ECOG-PS, UICC stage, CA 19-9, Prognostic CRP Classification, mGPS
44.	Wu et al <sup>104</sup> 2015	Prospective	Lung		China	366	NLR>2.68		Combination Chemotherapy	N/A	366	N/A	Multivariate: 1.778 (1.157-2.732) p=0.009	Metastasis, CRP
45.	Hong et al <sup>154</sup> 2015	Retrospective	Lung Cancer		China	919	NLR<5 Inverse: NLR>5		Chemotherapy and radiotherapy	N/A	892	N/A	Multivariate: 0.908 (0.721-1.144) p=0.413 Inverse Multivariate: 1.101 (0.874-1.387)	Stage, Response to treatment, LDH
46.	Yao et al <sup>103</sup> 2015	Retrospective	Prostate		Japan	57	NLR≥3.5		Docetaxel Chemotherapy	N/A	55	N/A	Multivariate: 2.728 (1.050-7.088) p=0.039	Biopsy Gleason Score, PSA value
47.	Wang et al <sup>129</sup> 2016	Retrospective	Cervical		China	60	NLR<2 Inverse: NLR>2		Cisplatin-based chemoradiotherapy	N/A	23	N/A	Multivariate: 0.268 (0.078-0.924) p=0.037 Inverse Multivariate: 3.731 (1.082-12.821)	Nil

48.	Beltran et al <sup>120</sup> 2016	Retrospective	T-cell lymphoma	Peru	83	NLR≥4	Combined Chemotherapy, Radiotherapy and Chemoradiotherapy	N/A	59	N/A	Multivariate: 4.73 (1.78-12.6) p<0.01	Performance Status
49.	Ferrucci et al <sup>142</sup> 2016	Prospective	Metastatic Melanoma	Italy	720	NLR≥3	Ipilimumab	N/A	662	N/A	Multivariate: 2.29 (1.86-2.82) p<0.0001	Sex, ECOG, Brain Mets, Liver Mets
50.	Zhang et al <sup>121</sup> 2016	Retrospective	RCC	China	373	NLR≥2.2	Combined Chemotherapy including Sorafenib and Sunitinib	N/A	373	N/A	Multivariate: 1.391 (1.022-1.894) p=0.036	Age, ECOG, IMDC Poor, Pathology, Fuhrman grade
51.	Kou et al <sup>108</sup> 2016	Retrospective	Pancreatic	Japan	306	NLR≥5	Combination chemotherapy with palliative intent	N/A	249	N/A	Multivariate: 2.54 (1.75-3.69) p<0.01	ECOG PS, Distant Metastasis, Initially unresectable, CEA, CA19-9
52.	Namikawa et al <sup>131</sup> 2016	Retrospective	Gastric	Japan	224	NLR≥4	Combination chemotherapy including trastuzumab	N/A	223	N/A	Multivariate: 1.651 (1.187-2.297) p=0.003	Histological type
53.	Moon et al <sup>131</sup> 2016	Prospective	Neck Squamous Cell Ca	Korea	153	NLR: Continuous	Combination chemotherapy and chemoradiotherapy	24	27	Multivariate: 4.13 (1.57-9.19) p=0.003	ECOG 1/0, BMI <18.5/others	
54.	Lee et al <sup>122</sup> 2016	Retrospective	Cholangiocarcinoma	Korea	221	NLR>5	Combination chemotherapy including Gemcitabine and 5-Fluorouracil based	N/A	197	N/A	Multivariate: 1.87 (1.33-2.62) p<0.001	Carcinoembryonic antigen, carbohydrate antigen 19-9, stage
55.	Ahn et al <sup>109</sup> 2016	Retrospective	Multiple Cancer Types	Korea	205	NLR≥10	Best supportive care	N/A	205	N/A	Multivariate: 1.54 (1.14-2.07) p=0.005	Cholangiocarcinoma, number of cycles of chemotherapy
56.	Choi et al <sup>134</sup> 2016	Retrospective	Pancreatic	Korea	396	NLR: 2.5-4.4 NLR: ≥4.5	Palliative Chemotherapy	N/A	396	N/A	Multivariate: 2.5-4.4: 1.659 (1.306-2.108) p<0.001 ≥4.5: 2.926 (2.181-3.927) p<0.001	ECOG PS, CA19-9

57.	Zaragoza et al <sup>144</sup> 2016	Retrospective	Melanoma	France	58	NLR week 1: continuous NLR week 1: ≥4	Chemotherapy including ipilimumab	N/A	22	N/A	Multivariate: Continuous: 1.10 (1.01-1.19) p=0.026 ≥4: 2.20 (1.01-4.78) p=0.047	LDH IU, Performance Status
58.	Li et al <sup>123</sup> 2016	Retrospective	Colorectal Ca Mets	China	110	NLR≤5 Inverse: NLR≥5	Combination chemotherapy including XELOX, FOLFOX and FOLFIRI	N/A	86	N/A	Multivariate: 0.99 (0.52-1.91) p=0.98 Inverse Multivariate: 1.01 (0.524-1.923)	Age, ALP Level, Ascites, PLR
59.	Hsieh et al <sup>193</sup> 2016	Retrospective	Gastric	Taiwan	256	NLR≤3	Combination Chemotherapy	N/A	248	N/A	Multivariate: 2.04 (1.22-3.40) p=0.007	Peritoneal Mets, NLR, mGPS, PG-SGA
No: LMR	Study	Type of Study	Cancer	Country	Patients (n)	Measure of SIR	Systemic Treatment	Cancer deaths (n)	Overall deaths (n)	Cancer survival (HR, 95%CI)	Overall survival (HR, 95%CI)	Independent Prognostic Factors
1.	Li et al <sup>124</sup> 2013	Retrospective	Nasopharyngeal	China	1547	LMR>5.220	Treatment with chemotherapy and radiotherapy	1457	1465	Multivariate: 0.669 (0.535-0.838) p=0.001 Inverse: 1.495 (1.193-1.869)	Multivariate: 0.558 (0.417-0.748) p=0.001 Inverse: 1.792 (1.337-2.398)	Sex, age, T stage, N stage, overall stage, treatment, prognostic measures.
2.	Rambaldi et al <sup>125</sup> 2013	Retrospective	B-Cell Lymphoma	Italy	700	LMR≤2.6	Systemic chemotherapy including rituximab	N/A	392 (10 Year survival)	N/A	Multivariate: 1.88 (1.32-2.70) p=0.001	IP>2
3.	Lin et al <sup>147</sup> 2014	Retrospective	SCLC	China	370	LMR≥4.56 Inverse: LMR≤4.56	Platinum based doublet chemotherapy	N/A	370	N/A	Multivariate: 0.530 (0.409-2.795) p<0.001 Inverse: 1.887 (0.358-2.445)	ALC, Histology, ECOG
4.	Lin et al <sup>148</sup> 2014	Retrospective	Nasopharyngeal	China	281	LMR≥5.07 Inverse: LMR≤5.07	Cisplatin based chemotherapy	N/A	255	N/A	Multivariate: 0.42 (0.30-0.59) p<0.001 Inverse: 2.381 (1.695-3.333)	Age, ALC

5.	Go et al <sup>136</sup> 2014	Retrospective	SCLC	Korea	188	LMR: Low	Platinum based chemotherapy	N/A	152	N/A	Multivariate: 1.472 (1.029-2.106) p=0.034	Stage
6.	Koh et al <sup>137</sup> 2015	Retrospective	Hodgkin Lymphoma	Korea	351	LMR<2.8	Active chemotherapy	38	48	N/A	Multivariate: 3.678 (1.008-13.41) p=0.049	LMR Only
7.	Jiang et al <sup>138</sup> 2015	Retrospective	Nasopharyngeal	China	672	LMR (≥2.475 vs. <2.475)	Active chemotherapy multiple modalities	N/A	458	N/A	Multivariate: 0.50 (0.41-0.60) p<0.001 Inverse: 2 (1.666-2.439)	N-stage, Number of metastatic lesions, Liver Mets
8.	Ho et al <sup>127</sup> 2015	Retrospective	Large B Cell Lymphoma	Taiwan	148	LMR<2.11	Standard R-chemotherapy.	N/A	41	N/A	Multivariate: 1.528 (0.751-3.111) p=0.242	Age, B-symptoms, ECOG, ALC, AMC, ALC/AMC PS
9.	Song et al <sup>189</sup> 2015	Retrospective	Colorectal	Korea	177	LMR≤3.4	Best supportive care and herbal therapy	N/A	177	N/A	Multivariate: 1.658 (1.092-2.518) p=0.018	mGPS, CA19-9, AST, KM treatment
10.	Simon et al <sup>1239</sup> 2016	Retrospective	Hodgkin's Lymphoma	Hungary	121	LMR≤2.11	Combination of chemotherapy and radiotherapy	N/A	13	N/A	Multivariate: 5.57 (1.53-20.25) p=0.003	PET 2 (positive)
11.	Lin et al <sup>149</sup> 2016	Retrospective	Metastatic Colorectal	China	488	LMR≥3.11 Inverse: LMR≤3.11	FOLFOX chemotherapy	N/A	479	N/A	Multivariate: 0.662 (0.501-0.875) p=0.004 Inverse: 1.511 (1.143-1.996)	Gender, ECOG Performance, Tumour differentiation, Pre-chemo AMC
No: PLR	Study	Type of Study	Cancer	Country	Patients (n)	Measure of SIR	Systemic Treatment	Cancer deaths (n)	Overall deaths (n)	Cancer survival (HR, 95%CI)	Overall survival (HR, 95%CI)	Independent Prognostic Factors
1.	Unal et al <sup>205</sup> 2013	Prospective	Lung	Turkey	94	PLR>194	Chemoradiotherapy including platinum based chemotherapy	N/A	81	N/A	Multivariate: 1.87 (1.20-2.91) p=0.006	Response to chemoradiotherapy, NLR

2.	Liu et al <sup>240</sup> 2013	Prospective	Lung	China	210	PLR≥152.6	First-line platinum-based chemotherapy	N/A	210	N/A	Multivariate: 2.025 (1.405-2.919) p<0.0001	Female sex, TNM stage IV, ECOG, p<0.0001
3.	Martin et al <sup>100</sup> 2015	Retrospective	Pancreatic	Australia	124	PLR≥200	Chemo for metastatic disease and radio for locally advanced	N/A	114	N/A	Multivariate: 1.58 (1.07-2.33) p=0.02	CA19-9, ALC, ANC, Platelet, NLR, mGPS, Alb, ECOG
4.	Li et al <sup>241</sup> 2015	Retrospective	HCC	China	243	PLR>111.23	Multiple Palliative Chemo	N/A	208	N/A	Univariate: 1.003 (1.002-1.004) p=0.002	White cell, Neutrophil, Platelets, NLR
5.	Jiang et al <sup>242</sup> 2015	Retrospective	Nasopharyngeal	China	1261	PLR ≥153.64	Chemo and Radio active	137	125	Multivariate: 1.84 (1.26-2.67) p=0.001	Multivariate: 1.83 (1.28-2.61) p=0.001	Age, Sex, Histology, TNM, EBV DNA
6.	Nakamura et al <sup>243</sup> 2015	Retrospective	Cervical	Japan	32	PLR>322.0	All patients treated with external radiotherapy and concurrent cisplatin based chemo	N/A	32	N/A	Multivariate: 4.204 (1.158-15.268) p=0.029	2 <sup>nd</sup> line chemotherapy, Pre-treatment
7.	Langsenhner et al <sup>244</sup> 2015	Retrospective	Prostate	Austria	374	PLP≥190	Radiotherapy	18	65	Multivariate: 3.99 (1.19-13.4) p=0.025	Multivariate: 1.87 (1.02-3.42) p=0.044	Neoadjuvant ADT, Secondary ADT, Gleason score ≥7,
8.	Cannon et al <sup>245</sup> 2015	Retrospective	Lung	USA	59	PLR>146	Stereotactic Radiation Therapy	N/A	28 (17 month follow up)	N/A	Multivariate: 4.0 (1.5-11.0) p = 0.006	PLR only
9.	Hong et al <sup>154</sup> 2015	Retrospective	Lung Cancer	China	919	PLR≥250	Chemotherapy and radiotherapy	N/A	892	N/A	Multivariate: 0.975 (0.783-1.215) p=0.824	Stage, Response to treatment, LDH
10.	Wu et al <sup>104</sup> 2015	Prospective	Lung	China	366	PLR>119.50	Combination Chemotherapy	N/A	366	N/A	Multivariate: 1.079 (0.729-1.596) p=0.705	Metastasis, NLR, CRP
11.	Kou et al <sup>108</sup> 2016	Retrospective	Pancreatic	Japan	306	PLR≥150	Combination chemotherapy with palliative intent	N/A	249	N/A	Multivariate: 0.96 (0.72-1.28) p=0.78	ECOG PS, Distant Metastasis, Initially unresectable, CEA, CA19-9, NLR

12.	Li et al <sup>23a</sup> 2016	China	110	PLR>162	Combination chemotherapy including XELOX, FOLFOX and FOLFIRI	N/A	86	N/A	Multivariate: 2.27 (1.32-4.03) p=0.003	Age, ALP Level, Ascites
No: Unassigned scores	Study	Country	Patients (n)	Measure of SIR	Systemic Treatment	Cancer deaths (n)	Overall deaths (n)	Cancer survival (HR, 95%CI)	Overall survival (HR, 95%CI)	Independent Prognostic Factors
1.	Yao et al <sup>20</sup> 2014	China	316	GAR>0.58	Active platinum based chemo	N/A	209	N/A	Multivariate: 1.65 (1.20-2.26) p=0.002	Albumin
2.	Zhou et al <sup>17</sup> 2015	China	367	CRP/Alb ratio (≥0.441)	Etoposide-based chemotherapy	N/A	258	N/A	Multivariate: 1.34 (1.04-1.73) p=0.025	Cancer stage, LDH level, PS
3.	Shibutani et al <sup>19</sup> 2015	Japan	66	AGR (>1.25)	Active Chemotherapy including platinum chemo	N/A	N/A (Only HR reported)	N/A	Multivariate: 2.247 (1.069-4.722) p=0.033	NLR
4.	Chan et al <sup>21</sup> 2015	Hong Kong	425	AAPR (>0.68)	Palliative chemo and radiotherapy	N/A	418	N/A	Multivariate: 2.185 (1.780-2.683) p<0.001	AJCC, BCLC, CLIP, CUPI, JIS
5.	Yamashita et al <sup>18</sup> 2016	Japan	79	CRP/Alb ratio (CAR) ≥7	Docetaxel-based chemotherapy	36	42	NA	Multivariate: 2.34 (0.91-6.05) p=0.08	ECOG PS≥1, PSA at docetaxel initiation, Hb≥12g/dL
6.	Zhou et al <sup>17</sup> 2016	China	276: Testing 379: Validated	CRP/Globulin Ratio ≥1.29	Chemotherapy including etoposide based regimens as well as cranial radiotherapy	N/A	Testing: 213 Validated: 205	N/A	Testing Multivariate: 1.35 (1.61-1.81) p=0.046 Validated Multivariate: 1.43 (1.05-1.95) p=0.022	ECOG-PS, Disease stage



**Title/Abstract reviewed for  
relevance n=9546**

**Excluded:**

- Curative surgery: n=3114
- Survival not primary measure: n=1225
- No full text: n=1195
- Infection (bacterial/viral): n=924
- Animal studies: n=2021
- Not English language: n=219
- Review: n= 149


**Full text review:  
n=699**

**Excluded:**


- Primary surgery: n=421
- PFS only: n=62
- Survival not expressed as HR (95%CI):  
n=47


**Studies included from reference  
list search: n=29**


**Included in systematic review: n=198**



Study or Subgroup	log[Hazard Ratio]	SE	Experimental Control		Weight	Hazard Ratio		Year	Hazard Ratio	
			Total	Total		IV, Random, 95% CI			IV, Random, 95% CI	
1.3.1 OS										
Viganó 2000	0.642	0.177	208	227	4.2%	1.90 [1.34, 2.69]	2000			
Axdorph 2000	0	0	0	0		Not estimable	2000			
Maréchal 2007	1.402	0.393	90	99	2.0%	4.06 [1.88, 8.78]	2007			
Lam 2007	0.052	0.016	57	145	5.6%	1.05 [1.02, 1.09]	2007			
Paralkar 2008	0.531	0.232	159	172	3.5%	1.70 [1.08, 2.68]	2008			
Ngo 2008	0.83	0.296	71	183	2.8%	2.29 [1.28, 4.10]	2008			
Shimoda 2010	1.967	0.966	83	83	0.5%	7.15 [1.08, 47.48]	2010			
Iwasa 2010	0.525	0.244	79	79	3.4%	1.69 [1.05, 2.73]	2010			
Yi 2011	0.531	0.229	278	278	3.5%	1.70 [1.09, 2.66]	2011			
Shim 2011	0.599	0.166	502	502	4.3%	1.82 [1.31, 2.52]	2011			
Trédan 2011	0.385	0.189	299	264	4.0%	1.47 [1.01, 2.13]	2011			
Lim 2012	0.748	0.353	49	50	2.3%	2.11 [1.06, 4.22]	2012			
Prakash 2012	0.859	0.296	314	486	2.8%	2.36 [1.32, 4.22]	2012			
Tsai 2014	0.683	0.343	479	522	2.4%	1.98 [1.01, 3.88]	2014			
Malik 2014	1.036	0.51	51	70	1.4%	2.82 [1.04, 7.66]	2014			
Imedio 2014	1.095	0.543	44	62	1.3%	2.99 [1.03, 8.66]	2014			
Stenman 2014	1.001	0.41	84	84	1.9%	2.72 [1.22, 6.08]	2014			
Kang 2014	0.693	0.341	168	168	2.4%	2.00 [1.02, 3.90]	2014			
Ulas 2014	0.247	0.136	391	462	4.7%	1.28 [0.98, 1.67]	2014			
Narwani 2015	2.235	0.564	22	38	1.2%	9.35 [3.09, 28.23]	2015			
Koo 2015	0.281	0.041	3494	3888	5.5%	1.32 [1.22, 1.44]	2015			
Go 2015	0.651	0.298	0	134	2.8%	1.92 [1.07, 3.44]	2015			
Xue-Feng 2015	0.075	0.285	127	127	2.9%	1.08 [0.62, 1.88]	2015			
Martin 2015	0.753	0.215	114	124	3.7%	2.12 [1.39, 3.24]	2015			
Helissey 2015	2.407	0.573	26	56	1.2%	11.10 [3.61, 34.13]	2015			
Kao 2015	0.737	0.261	69	143	3.2%	2.09 [1.25, 3.49]	2015			
Wild 2015	1.277	0.342	88	101	2.4%	3.59 [1.83, 7.01]	2015			
Choi 2016	0.322	0.117	396	396	4.9%	1.38 [1.10, 1.74]	2016			
Uemura 2016	1.329	0.569	22	22	1.2%	3.78 [1.24, 11.52]	2016			
Kou 2016	-0.223	0.156	249	306	4.4%	0.80 [0.59, 1.09]	2016			
Dorajoo 2016	0.259	0.112	480	482	4.9%	1.30 [1.04, 1.61]	2016			
Subtotal (95% CI)			8493	9753	91.4%	1.77 [1.54, 2.03]				
Heterogeneity: Tau² = 0.08; Chi² = 181.16, df = 29 (P < 0.00001); I² = 84%										
Test for overall effect: Z = 8.05 (P < 0.00001)										
1.3.2 CSS										
Axdorph 2000	0.94	0.455	48	145	1.7%	2.56 [1.05, 6.25]	2000			
Al Murri 2006	1.227	0.48	51	96	1.6%	3.41 [1.33, 8.74]	2006			
Ramsey 2007	0.967	0.33	102	119	2.5%	2.63 [1.38, 5.02]	2007			
Uemura 2016	1.329	0.569	22	22	1.2%	3.78 [1.24, 11.52]	2016			
Moon 2016	1.335	0.451	24	153	1.7%	3.80 [1.57, 9.20]	2016			
Subtotal (95% CI)			247	535	8.6%	3.05 [2.08, 4.47]				
Heterogeneity: Tau² = 0.00; Chi² = 0.78, df = 4 (P = 0.94); I² = 0%										
Test for overall effect: Z = 5.73 (P < 0.00001)										
Total (95% CI)										
			8740	10288	100.0%	1.87 [1.63, 2.15]				
Heterogeneity: Tau² = 0.09; Chi² = 207.73, df = 34 (P < 0.00001); I² = 84%										
Test for overall effect: Z = 8.93 (P < 0.00001)										
Test for subgroup differences: Chi² = 6.93, df = 1 (P = 0.008), I² = 85.6%										
0.01 0.1 1 10 100										
Favours [experimental] Favours [control]										







Study or Subgroup	log[Hazard Ratio]	SE	Experimental Total	Control Total	Weight	Hazard Ratio IV, Random, 95% CI	Year	Hazard Ratio IV, Random, 95% CI
<b>1.10.1 OS</b>								
Lin 2014(ii)	0.868	0.172	255	281	9.4%	2.38 [1.70, 3.34]		
Rambaldi 2013	0.631	0.183	392	700	8.5%	1.88 [1.31, 2.69]	2013	
Li 2013	0.583	0.149	1465	1547	11.7%	1.79 [1.34, 2.40]	2013	
Go 2014	0.387	0.183	152	188	8.5%	1.47 [1.03, 2.11]	2014	
Lin 2014(i)	0.635	0.49	370	370	1.4%	1.89 [0.72, 4.93]	2014	
Song 2015	0.506	0.213	177	177	6.6%	1.66 [1.09, 2.52]	2015	
Koh 2015	1.302	0.66	48	351	0.8%	3.68 [1.01, 13.40]	2015	
Ho 2015	0.424	0.363	41	148	2.5%	1.53 [0.75, 3.11]	2015	
Jiang 2015	0.693	0.097	458	672	20.6%	2.00 [1.65, 2.42]	2015	
Simon 2016	1.717	0.659	13	121	0.8%	5.57 [1.53, 20.26]	2016	
Lin 2016	0.413	0.142	479	488	12.6%	1.51 [1.14, 2.00]	2016	
<b>Subtotal (95% CI)</b>			<b>3850</b>	<b>5043</b>	<b>83.2%</b>	<b>1.84 [1.64, 2.07]</b>		
Heterogeneity: $\tau^2 = 0.00$ ; $\chi^2 = 10.86$ , $df = 10$ ( $P = 0.37$ ); $I^2 = 8\%$								
Test for overall effect: $Z = 10.29$ ( $P < 0.00001$ )								
<b>1.10.2 CSS</b>								
Li 2013	0.402	0.115	1457	1547	16.8%	1.49 [1.19, 1.87]	2013	
<b>Subtotal (95% CI)</b>			<b>1457</b>	<b>1547</b>	<b>16.8%</b>	<b>1.49 [1.19, 1.87]</b>		
Heterogeneity: Not applicable								
Test for overall effect: $Z = 3.50$ ( $P = 0.0005$ )								
<b>Total (95% CI)</b>			<b>5307</b>	<b>6590</b>	<b>100.0%</b>	<b>1.78 [1.58, 1.99]</b>		
Heterogeneity: $\tau^2 = 0.01$ ; $\chi^2 = 13.61$ , $df = 11$ ( $P = 0.26$ ); $I^2 = 19\%$								
Test for overall effect: $Z = 9.74$ ( $P < 0.00001$ )								
Test for subgroup differences: $\chi^2 = 2.60$ , $df = 1$ ( $P = 0.11$ ), $I^2 = 61.5\%$								

